

# Range Nutrition

The key to range nutrition is matching the needs of the grazing animal with the available forage. The needs of the animal will vary with the age of the animal, if mature; stage of pregnancy, if lactating; and the age of nursing offspring. The nutritional value of forage will vary with the stage of plant development and environmental conditions.

This section is modified from a publication by C. Wayne Cook, which was accessed through the Colorado State University website on May 27, 2005.

**TIP: The quality of a range plant is judged primarily on palatability, or how readily animals eat it, and its nutritive content with respect to its phenological development. Some plant species are eaten only during early growth and, in special cases, only certain portions of a particular species are consumed during the latter stages of development.**

During their annual life cycles, individual plant species vary in nutrient content. Most forage plants are high in nutrients during early growth. However, as they mature, they lose nutrients markedly. Some plant species mature rather rapidly and, as a result, decrease substantially in nutritional value; other species mature rather slowly and, consequently, remain high in nutrient content during an extended period. Still other plant species, even after they mature, appear to cure rather well and retain comparatively high quantities of nutrients for indefinite periods.

The true test of the nutrient value of a forage species, or a mixture of species, is the ability of the usable forage to meet the nutritional requirements for the physiological functions being performed by the grazing animal during the various seasons of the year. Most physiological functions of the grazing animal are performed as everyday functions and generally are referred to as "maintenance requirements," which include travel, mastication and digestion, maintaining body temperature and growing hair or wool.

The nutrient evaluation of most range areas can be based on how much protein, energy and phosphorus are in the forage plants. Phosphorus, digestible protein and energy content are the three main nutrients that are most important in evaluating the status of range nutrition.

The comparative nutritive value among the forage classes (browse, forbs and grasses) can be discussed best by measuring their apparent ability to meet the nutritional requirements of large herbivores for the more important physiological functions such as maintenance, gestation, growth and lactation.

In addition to the influence of relative preference and floristic composition on nutritive content of range forage, factors such as stage of growth and variable site conditions also are important.

## Nutritional Content and Stage of Growth

For the most part, rangelands of the West can be evaluated for nutritional value on the basis of phosphorus, digestible protein and digestible energy. Some local areas also may have other nutrients that should be considered. These nutrients in the Dakotas, Minnesota and south-central Canada include copper and zinc.

During the period of initial growth, and for a time thereafter, all forage classes are high in nutrient content, and their nutritional contents are not materially different. However, as growth stages advance, the nutritional differences among forage classes become more evident. Researchers generally agree that young, growing animals and lactating animals have comparable requirements when expressed on a percentage basis.

The common belief is that animals grazing green plant growth on spring and summer ranges receive adequate nutrients; however, on fall and winter ranges where plants are dry and mature, the diets of grazing animals are thought to be deficient in some nutrients.

Several studies have shown that chemical composition varies with season, largely as a result of changes in the stem-to-leaf ratio and the normal maturing process that causes a translocation of nutrients in the plant parts. In addition to the actual decrease in chemical content of herbage with advancement of growth, the forage likewise decreases in digestibility because of lignification and calcification of the plant material.

### Digestible Protein

The correlation between total protein and digestible protein in the mixed ration normally used in farmstead feeding is rather high. In range forages, the digestibility of protein may vary from 70 percent in early growth to as low as 10 to 15 percent in the quiescent stages. Thus, evaluating grazing animals' diet on the basis of digestible protein rather than on total protein appears to be more logical unless standards have considered the high variability in the digestibility of protein as plants mature.

Grass species decline in digestible protein rather rapidly and generally fail to meet the lactation requirements at about the time they come into full anthesis (flowering) (Figure 6). This is the result of a rather rapid loss of total protein and a more rapid decrease in digestibility of protein that occurs with advanced growth. Grasses, in general, lose 75 percent of their protein during the period from early growth to seed formation; on the other

hand, browse lose only about 40 percent of their protein content during a similar period. As a result, grasses that have matured are considered poor sources of digestible protein. Shrubs, however, are considered good sources of digestible protein during most of their active growing period; and even after they reach full maturity, they continue to meet gestation requirements.




Figure 6. Average content of digestible protein for the three forage classes compared with phenological development and herbivore requirements for lactation and gestation.

Forbs are intermediate between browse and grass with respect to protein content during most seasons. Most forb species fail to furnish adequate digestible protein to meet the requirements of animal gestation after reaching the fruiting stage.

### Energy

In some cases, neither grass energy nor digestible energy in shrubs is considered a good index to the true energy values of forage because of the high content of essential oils, resins or waxes that suggest high energy content, but these materials are not available for livestock metabolism. The digestible energy values for browse shown in Figure 7 have been adjusted for species high in essential oils so that the trends are based on realistic digestible energy values.

Shrubs are not considered good sources of energy after they reach the phenological stage of fruit development. Thereafter, they generally fail to meet the energy requirement for animals in gestation.

Grasses generally are considered good or excellent sources of energy, primarily because of their high content of cellulose. Forbs are intermediate between shrubs and grasses in energy-furnishing constituents and, like shrubs; they generally fail to meet the energy requirements for gestation after reaching full maturity and dormancy.




Figure 7. Average content of digestible energy for three forage classes compared with various stages of phenological development and requirements for lactation and gestation.

## Phosphorus

Even when mature, shrubs generally are considered good sources of phosphorus for general animal maintenance and gestation unless they are deciduous. Even deciduous shrubs are perhaps only borderline if the young twigs are eaten readily (Figure 8). Most forbs have a phosphorus content that is only slightly lower than that of shrubs.

Grasses, however, are low in phosphorus soon after they form seed, so they are considered poor sources of phosphorus when mature and not actively growing. Most grasses lose considerable phosphorus content when temporarily forced into dormancy by even brief periods of drought. However, when precipitation occurs and growth is renewed, the phosphorus content increases and lactation requirements again are met.




Figure 8. Average percent of phosphorus for three forage classes compared with stage of growth and herbivore requirements for lactation and gestation.

**TIP:** Maintaining or improving the diversity of grasses, forbs and shrubs in your rangeland pastures will permit livestock the opportunity to select the forages that will best meet their nutritional needs throughout the grazing season.

## Copper and Zinc

Range and pastureland in the northern Great Plains region are considered deficient especially of copper but also of zinc by the seed head stage of development. Cattle will become deficient of copper by early summer and may show signs of deficiency by midsummer, especially red pigmented cattle when the water source is high in sodium. If the drinking water is known to be high in sodium, adding extra copper to the mineral program is highly recommended starting midsummer and through the winter feeding program.

Sheep can be poisoned easily by excess copper supplementation, so caution is warranted when adding copper to the pasture when sheep are present. Year-to-year variation in the zinc content of pasture plants occurs; however, it becomes borderline to deficient by midsummer.

## ■ Nutritive Value as Affected by Site

Site conditions are important because they influence the growth characteristics of range plants and, thus, indirectly affect their nutritive value. Sites also indirectly affect the chemical content of plants and plant parts through soil and plant development, water runoff, intensity of shade and other environmental factors.

Shrubs and forbs, when they approach maturity, are considerably leafier on less favorable growing sites. Grasses in advanced stages of growth are, likewise, leafier on poorer sites than on more favorable sites, but differences between sites are less conspicuous for grasses than for shrubs or forbs. As a result, plants on unfavorable sites are more palatable, more digestible and, therefore, more nutritious than plants on the favorable sites.

Nutrient content of plants on poor sites during advanced growth stages generally are considered more nutritious because of the finer material and the presence of more leafy material, compared with stems. The differences in stem-to-leaf ratios would, to a large degree, account for chemical differences between plants growing on favorable and unfavorable sites because leaves are higher in ether extract, protein, ash, calcium, phosphorus, and nitrogen-free-extract, whereas stems are higher in lignin, crude fiber and cellulose.

Predominantly grass ranges such as mixed and tall grasses are considered deficient in phosphorus, protein and vitamin A when the plants approach hard seed formation or maturity (See Figures 6, 7 and 8). Therefore, grass ranges generally are believed to be deficient in these three nutrients in late summer, fall and winter, and supplementation should be considered.

While generally not considered a nutrient, water is critical to livestock performance. Forage intake is directly proportional to water intake, and providing adequate, high-quality water is important. Refer to Tab 19 for further information about water developments.

### Other References

1. Grazing Land Animal Nutrition Lab at Texas A&M University available at: <http://cnrit.tamu.edu/ganlab/>
2. Livestock nutrition publications: <http://cnrit.tamu.edu/ganlab/>
3. Grasses for the Northern Plains: Vol. I Cool-season. R1323 – North Dakota State University Extension Service and USDA Plant Material Center ([www.nrcs.usda.gov/Internet/FSE\\_PLANTMATERIALS/publications/ndpmcbk7681.pdf](http://www.nrcs.usda.gov/Internet/FSE_PLANTMATERIALS/publications/ndpmcbk7681.pdf)). Grasses for the Northern Plains, Volume 1 - Cool-Season
4. Grasses for the Northern Plains: Vol. II Warm-season. R1390 – North Dakota State University Extension Service and USDA Plant Material Center ([www.nrcs.usda.gov/Internet/FSE\\_PLANTMATERIALS/publications/ndpmcbk9425.pdf](http://www.nrcs.usda.gov/Internet/FSE_PLANTMATERIALS/publications/ndpmcbk9425.pdf)). Grasses for the Northern Plains, Volume 2 - Warm-Season

# Factors Affecting the Nutritive Value of Range Forage

KERMIT OELBERG

Senior Student, Department of Forestry and Range Management, Washington State College, Pullman, Washington

Antoine Lavoisier (1743-1794), a great French chemist, is accredited as being the founder of the science of nutrition. His famous respiration experiments led him to state "La vie est une fonction chimique" or "Life is a function of chemistry." Since Lavoisier's time many advancements have been made in the field of animal nutrition.

The nutritive value of any forage is dependent upon its content of energy-producing nutrients as well as its content of nutrients essential to the body, namely, protein, minerals and vitamins.

Forages supply energy mostly in the form of carbohydrates. The carbohydrate fraction makes up from 60 to 80 percent of the dry matter. The higher carbohydrates may be broken up into three classes: (1) starch, readily and nearly completely digested by the animal; (2) cellulose, digested mostly by bacteria in the rumen of ruminants; and (3) hemicelluloses, intermediate between starches and cellulose and broken down by weak acids and alkalis. In chemical analyses, these higher carbohydrates are arbitrarily grouped into two classes, crude fiber and nitrogen-free extract. Crude fiber consists almost entirely of cellulose and lignin together with some resistant hemicellulose. Lignin is essentially non-digestible by most domestic animals. However, Utah studies have shown that deer are able to digest up to 42 percent of this constituent in birchleaf mahogany (*Cercocarpus montanus*) (Smith, *et al.*, 1956). Nitrogen-free extract consists mainly of starch and most of the hemicelluloses.

Protein commonly means nitrogen multiplied by the factor 6.25;

however, 25 to 50 percent of the total nitrogen may be non-protein, essentially the amides and amino acids (Maynard, 1937). The value of protein is dependent principally upon its supply of certain amino acids which are required for the formation of body protein and which cannot be manufactured by the body. The amount of digestible protein furnished by a plant depends upon the plant species and the class of livestock using the plant. In addition, digestibility of protein varies appreciably between domestic animal species and somewhat between individuals of a given species.

Minerals are essential for the proper functioning of the bodily processes. Aside from sodium, chlorine, calcium, and phosphorus, most of the essential elements are ample in forages except when grown on mineral-deficient areas, i.e., cobalt in Florida.

The precursors of vitamin A and vitamin D are frequently limiting in range forage. Carotene is the precursor of vitamin A, which is manufactured in the animal body, and ergosterol is the precursor of vitamin D in plants. Cholesterol is the animal form of vitamin D precursor and like ergosterol, it needs sun exposure for transformation into vitamin D (Maynard, 1947). Both of these vitamins play an important role in body processes and growth.

## Determination of Nutritive Value

Precise information relating to the feeding value of range forages is notably lacking. This is a reflection of the complexity of plant analysis, as well as the difficulty of interpreting results in terms of actual feeding values.

Early investigators resorted to

chemical analyses of various species, and through the years developed a fairly reliable system of chemical analysis procedures. However, it was noted that chemical analysis data had to be interpreted with care. Many plants, found to be highly nutritious by chemical analysis, proved to be worthless as animal forage because they lacked palatability.

Most students have been concerned with only the major forage plants, chemical analyses being made on the parts thought to be consumed or bulk samples of plant material (Clarke and Tisdale, 1945; Hart, *et al.*, 1932; Stoddart and Greaves, 1942). The chemical analysis of consumed plant forage is in itself an incomplete measure of nutritive value, but it can be used as a guide when interpreted with the results of digestibility trials that have been conducted with similar forages (Cook and Harris, 1950).

Later experimenters used digestion trials to test nutritive value. For obvious reasons, the data secured from these trials were not indicative of the actual grazing habits of animals, which show forage preferences. Forages also vary as to palatability and composition when artificially harvested and fed to livestock by man. Consequently, some present-day workers have endeavored to improve field digestion trials by use of specially designed fecal and urinal bags attached to the grazing animal. Even this advanced procedure has its weaknesses, and will require many more years of refinement through trial before the accuracy of results approaches an acceptable level.

## Factors Influencing Nutritive Value

The factors influencing the nutritive value of range forage are many and the degree to which they are interrelated may vary considerably from one area to another. The study of these influences is indeed complex, for the experimenter must attempt to study individual factors while keeping the remaining ones as nearly constant as possible.

sible. It is essential that due credit be attributed to the correct factor or complex of factors.

The nutritive value of range forage is influenced in a major way by: (1) stage of maturity, (2) edaphic influences, (3) plant species, (4) climate, (5) animal class, and (6) range condition. In this report, an attempt will be made to show how these important factors influence the nutritive value of forage.

#### Stage of Maturity

The stage of growth seems to be the most important factor affecting the chemical composition and digestibility of range forage. In general all forages are highly succulent in early growth, which markedly enhances their palatability. In addition, their high protein content in relation to a low fiber content at this stage makes them highly nutritious as livestock forage. Thus, grasses and forbs are referred to as "watered concentrates" while in the early stages of growth.

Striking nutritive differences exist between forage classes (browse, grasses and forbs) at maturity. Browse species are less affected by summer drought periods than are forbs and grasses because of their deeper root systems.

#### Browse:

Protein content of browse tends to decrease as the season advances except in certain species which show an increase in protein during the moist fall period. In snowberry (*Symphoricarpos rotundifolius*), protein decreased from an average of 11.24 percent on July 15 to 9.19 on August 15 to 6.58 percent on September 15 (Stoddart, 1941). However, when various plant parts of *Symphoricarpos vaccinooides* were analyzed separately, it was found in some cases that the protein content increased with advanced maturity (Cook and Harris, 1950a).

Little change occurred in the nutritive value of winter range plants in Utah during the winter grazing season (November 18 to March 14). Normal leaching and translocation of nutrients occurred

prior to the winter grazing season, accounting for the constancy in chemical composition of these winter species (Cook and Harris, 1950).

The trend in crude fiber content in regard to stage of maturity is normally the reverse of protein. In California, Gordon and Sampson (1939) found that half-shrubs, *Lotus scoparius* and *Lupinus albus*, had an increasing amount of crude fiber until late summer when it decreased due to a larger proportion of leafage on the shoots. In deciduous shrubs such as *Aesculus californica* and *Sambucus glauca*, the crude fiber content stayed rather uniformly low at all stages of growth. As the percentage of crude fiber increases, digestibility usually decreases because crude fiber is resistant to decomposition and it often envelops digestible nutrients rendering them unavailable (Maynard, 1937).

"Nitrogen-free extract" (mainly starches and hemicelluloses) normally decreases with advancing season. Cook and Harris (1950a) found that nitrogen-free-extract content decreased in all plant parts of snowberry.

Phosphorus content normally parallels that of protein in regard to seasonal changes. Phosphorus and magnesium both decreased significantly in *Symphoricarpos rotundifolius* with advancing season (Stoddart, 1941). Deciduous shrubs in California also showed a marked decrease in phosphorus content toward maturity (Gordon and Sampson, 1939).

Calcium, in contrast, generally increases as the season advances (Savage, et al., 1947). The calcium content of several deciduous shrubs was observed to increase uniformly from early leaf stage to maturity (Gordon and Sampson, 1939). The fact that calcium content increased with maturity was explained on the basis of the increased amount of cellular material which is composed principally of this element. McCreary (1927) has suggested that the late-season increases in calcium and ash may be

attributed to dust accumulations. Hart, et al., (1932) report no seasonal trend in calcium content. These reported differences in results seem to indicate that there are several interrelated and poorly understood factors which influence the calcium content of shrubs.

Vitamins, essential for animal metabolism, also vary with season. Vitamins are unstable in dry forage and quickly disintegrate as leaves and stems desiccate. Carotene content of shrubs decreases, although slower than in grasses and forbs, as the season progresses.

#### Forbs:

Generally speaking, the leaves and blades of grasses and forbs mature earlier than shrubs. Seasonal changes alter the nutritive value of forbs (especially spring forbs) much faster than browse or most grass species because the succulent tissues of forbs are subject to desiccation upon exposure to the high temperatures and long photoperiods of the summer months. Therefore, the period of high nutritive value of most forbs is restricted to the early portion of the growing season. The rate of decrease in protein and crude fiber in the broadleaf herbs except legumes is most rapid from early leaf stage to the late blossom period. In California legumes studied by Gordon and Sampson (1939), the protein content declined slowly and uniformly throughout the season, the relatively high content at maturity resulting in part from the high protein level of the seeds of many species. Cook and Harris (1950a) reported an orderly decrease in protein content of yarrow (*Achillea lanulosa*) except for a slight increase at the end of the growing season, attributed to improved moisture conditions during this period.

In general, phosphorus content parallels that of protein in most forbs. Calcium was found to increase with maturity in all plants and plant parts studied by Cook and Harris (1950a). Several species of *Erodium* were found to be remarkably high in silica-free ash

and calcium at maturity (Gordon and Sampson, 1939). Silicon, iron and aluminum are "hard" elements (not essential to animals); consequently, plant species having a minimum of these elements are most desirable.

#### Grasses:

Seasonal changes affect grasses in much the same way as forbs except that certain perennial grasses retain their nutrients after maturity.

Native and tame grasses of western North Dakota lost on an average 71 percent of their protein by September 30 (Whitman, *et al.*, 1951). In an extensive study of forage plants in Utah by Cook and Harris (1950), the grasses had an average protein content of 8.2 percent in early season (July 10 to August 4), 7.2 percent in mid-season (August 15-24) and 4.5 percent in the late season (August 25 to September 13). *Poa scabrella*, a perennial grass, had a protein content of approximately 20 percent at early leaf stage and only 5 percent at maturity; the annuals, *Bromus mollis* and *Avena fatua*, dropped 89 percent from early leaf stage to maturity (Gordon and Sampson, 1939).

Native range grasses of western North Dakota were seen to increase in crude fiber and decrease in phosphorus and carotene with advancing maturity. By the end of September, these native grasses had lost 87 percent of their carotene and 66 percent of their phosphorus (Whitman, *et al.*, 1951). Studies of pasture grasses have shown phosphorus to vary directly with crude protein content. Calcium, on the other hand, generally increases in grass tissue with advancing season. In Montana grasses, lignin increased from 5 percent in May to 18 percent in September (Patton and Gieseke, 1942).

#### Edaphic Influences

The physical and chemical properties of soils exert almost unlimited influence upon the nutrient content of plants. Orr (1929) concluded that the mineral composition within a species was deter-

mined primarily by soil as shown by the response to fertilizers. It has long been known that plants grown on soils rich in certain nutrients usually are also rich in these nutrients.

Physical properties of soil such as texture and porosity affect the nutritive quality of forage more or less indirectly. Poorly aerated soils greatly limit or decrease the absorption of essential elements, especially phosphorus. Soils rich in biotic life show enhanced aeration and fertility.

Chemical properties of the soil may determine the nutrients that plants are able to absorb. For example, phosphorus is most available between pH 6 and pH 7. Phosphorus in soils of low pH reacts chemically with hydrous oxides of iron, aluminum and magnesium to form insoluble compounds which are unavailable to plants. At pH 7 and above, phosphorus again becomes insoluble as calcium phosphate. Holtz (1930) showed that the phosphorus content of oats followed the *total* phosphorus content of the soil, whereas that of red clover followed the *available* phosphorus content of the soil.

Range plants growing on limestone outcroppings in California had relatively high calcium content (Hart, *et al.*, 1932).

Daniel (1934) found that certain species were normally high in calcium and phosphorus even when grown on soils relatively low in these minerals, and plants normally low in calcium and phosphorus had low contents of these minerals when grown on rich soils. This suggests genetic characteristics which are inherent to the specific plants.

Burning of heavy vegetation modifies both the physical and chemical and biological properties of the soil. Mild burning of a range area generally results in the release of minerals, making them available for plant absorption. Wild oats in California (24) had increased percentages of protein, nitrogen-free extracts, crude fat, calcium and

phosphorus the first year following a burn (Hart, *et al.*, 1932).

#### Climatic Influences

Climatic factors such as temperature, humidity, precipitation, light intensity and altitude may be dominant in controlling the nutritive value of plants. Although plants are dependent upon the soil for their mineral nutrients, climatic factors affect respiration, assimilation, photosynthesis and metabolism to the extent that the mineral and organic matter content of plants may be strongly modified by climatic factors even though grown on the same soil.

Plants of *Bromus inermis* grown under full sunlight were found by Watkins (1940) to have higher carbohydrate and lower protein content than plants grown in shade.

Precipitation may have direct and indirect influences upon the quality of forage plants. Rainfall, in general, tends to increase nitrogen, phosphorus and ether extract (the soluble fat constituent). Oklahoma studies by Daniel and Harper (1934) revealed a close relationship between precipitation and the calcium and phosphorus balance in *Andropogon scoparius*. Increased precipitation resulted in an increase in phosphorus and a decrease in calcium and vice versa. Watkins (1940) has shown that droughts may decrease the phosphorus and protein contents and increase the calcium and crude fiber contents. However, Ferguson (1931) found that both calcium and phosphorus decreased in forage during periods of drought. Scott (1929) found no significant relationship between precipitation and the calcium-phosphorus content of native forage species.

Precipitation greatly alters plant nutrients when they are matured. Leaching causes a decrease in all plant nutrients except crude fiber; thus the plant forage becomes less digestible (Guilbert, *et al.*, 1931).

Temperature seems to be the most important factor governing phenology. Low temperatures tend to initiate the transformation of starches into plant sugars which

are used in plant metabolism. Benedict (1940) found that low night temperature increased the dry weight of *Agropyron smithii* and *Andropogon furcatus* when subjected to both long and short days.

Altitude affects plant composition through the interrelation of factors such as light intensity, carbon dioxide concentration and precipitation. Western forage plants generally increase in feeding value with increasing altitude (Roberts, 1926). The nitrogen content of high altitude plants seems to be higher than that of plants grown at lower elevations (McCreary, 1927).

#### Plant Species

A plant may have satisfactory amounts of inorganic and organic nutrients as forage for livestock but is of little value if it lacks palatability. Big sagebrush (*Artemisia tridentata*) is an example. Therefore digestion experiments are mandatory when determining the real forage value of plant species. Furthermore, the palatability and abundance of the various species determine the botanical composition of the grazing animal's diet.

Many workers have concluded that the plant species is more important than the soil or management practices in determining the mineral composition of the forage. However, the soil may govern the type of plants which grow upon an area. Legumes contain more calcium than grasses and their calcium-phosphorus ratio is higher. Leguminous species carry their own nitrogen-factory with them and are able to build up nitrate reserves in the soil which may become available to associated forage species.

Browse species are generally higher in protein content than are grasses and forbs. Browse species retain their protein, carotene and carbohydrates better during periods of drought than either grasses or forbs because of their deep root systems and ability to store food reserves in their stems (Stoddart and Smith, 1955). Forbs other than leguminous species tend to be slightly higher in protein, phos-

phorus and calcium than grasses during their growth period; however, most of them succumb to the elements soon after maturity.

On winter range areas, browse furnishes the major components of the grazing animal diet because the nutrients in the grasses on these winter ranges are able to supply large quantities of metabolizable energy. Browse species are relatively low in energy values because of their high lignin content in comparison to their cellulose content (Cook and Harris, 1950).

#### Livestock Class

The various classes of livestock exhibit different behavior patterns when grazing. Sheep and goats tend to be more selective in their grazing, eating plant parts more often than the whole. Minor food preferences also exist among members of the same species.

Non-ruminants such as swine can make only limited use of range forage due to their limited digestive capacity. In certain sections of the United States, hogs are allowed to run at large, their diet consisting largely of mast (acorns and nuts) and limited forage. Hogs utilize such concentrated feeds efficiently, but cannot make good use of sizeable quantities of roughage.

Horses prefer grasses to all other forages. Although they are monogastric animals, the presence of a caecum or "blind gut" in their digestive systems enables them to digest large quantities of roughage.

Ruminants are physiologically adapted for the digestion of complex cellulose and to a limited extent lignin. Micro-organisms within the rumen act to break these complex substances into simpler carbohydrates which the ruminant can digest and metabolize.

Ruminants may eat some plants more readily than others, without relationship to the digestibility or nutritive value of these plants. Many palatable plants are actually very low in nutritive value.

Cattle, like horses, prefer grasses to browse or forbs. Cattle will eat mature, hardened forages more readily than sheep will. Horses

and cattle also prefer certain grass species to others.

Sheep prefer forbs and tender grasses, whereas goats prefer browse species. Sheep will graze *Bromus tectorum* when young and succulent but avoid it when it reaches maturity. Grass species such as *Poa secunda* which tend to be rather wiry are avoided by sheep.

Due to these forage preferences, many range areas are well suited for grazing by two classes of livestock or alternate grazing by each class in successive years.

#### Range Condition

Range condition is influenced essentially by the interplay of grazing intensity, season of use, grazing class, soil and climatic influences.

Grazing intensity influences the ultimate nutrient quality of forage. Livestock normally consume the leaves and more tender stems first and reject the fibrous plant parts. This reduces the photosynthetic area of the plant and the root-shoot balance is disturbed (Cook, *et al.*, 1948). These Utah studies showed that available protein, phosphorus, cellulose and metabolizable energy in the forage decreased with heavier utilization while the lignin content increased. Digestibility of protein and cellulose was found to decrease 10 and 67 percent respectively in some cases as utilization increased.

Native range grasses in Kansas clipped frequently yielded less forage but of a higher protein content than if clipped less often (Aldous, 1926). Sampson and Malmsten (1926) found that the herbage of *Agropyron violaceum* and *Bromus polyanthus* clipped four times a season contained less crude fiber and nitrogen-free extract, more crude protein and approximately the same amount of either extract than forage clipped only once per season. Immature herbage seems to lack "substance." Crude protein in more developed herbage exists in the form of amino acids which are completely assimilated by animals; whereas, the protein of young, succulent forage occurs in the amide

form and is not assimilated by the animal to a very high degree.

The stage of succession is also an important part of range condition. In general, climax vegetation of grassland areas produces the greatest quantity of desirable dry matter possible on a given site without irrigation or fertilization. Although the introduced cheatgrass (*Bromus tectorum*) yielded as much forage per acre as crested wheatgrass or native bunch grasses, its period of high nutritive value was limited to a short period during the spring months (Hull and Pechanec, 1947).

### Summary

The nutritive value of range forages is affected by stage of maturity, edaphic conditions, climatic influences, plant species, animal class and range condition.

Stage of maturity seems to influence forage quality more than any other factor. Protein, nitrogen-free extract, ether extract, carotene and phosphorus tend to decrease with advancing maturity, whereas crude fiber, lignin and calcium increase. The trend is more abrupt in forbs than in browse species, and intermediate in grasses.

The physical and biological properties of the soil, aeration, texture, and biotic influences are important in regulating physiological processes in plants. Chemical properties of soil, such as pH, available minerals and fertility, control to a certain extent the absorption of minerals by plants. Burning may modify both chemical and physical characteristics of soil. Generally, burning releases many unavailable elements.

Climate affects forage value considerably. Increased precipitation tends to increase the phosphorus content and decrease the calcium content of plants. Light intensity, temperature and carbon dioxide concentration limit forage value if they occur in sub-optimum quantities.

Plant species vary greatly as to palatability, digestibility and nutritive composition.

Animals possess certain forage preferences which increase the problem of proper grazing practices. Range condition is affected mostly by grazing intensity as it operates through other factors. In general browse and perennial grasses withstand grazing better than forbs. The most palatable species are most seriously affected by excessive grazing.

Grazing intensity on a given area should be ample to remove the maximum amount of forage without detriment to the potentialities of the site.

### ACKNOWLEDGMENT

The author is grateful to Professor GRANT A. HARRIS of the Department of Forestry and Range Management, Washington State College, for his assistance and interest in this problem.

### LITERATURE CITED

- ALDOUS, A. E. 1930. Effect of different clipping treatments on the yield and vigor of prairie grass vegetation. *Ecology* 11(4): 752-759.
- BENEDICT, H. M. 1940. Effect of day length and temperature on the flowering and growth of four species of grasses. *Jour. Agr. Res.* 61(9): 661-671.
- CLARKE, S. E. AND E. W. TISDALE. 1945. The chemical composition of native forage plants of southern Alberta and Saskatchewan in relation to grazing practices. *Canada Dept. Agr. Tech. Bull.* 54.
- COOK, C. W. AND L. E. HARRIS. 1950. The nutritive content of the grazing sheep's diet on summer and winter ranges of Utah. *Utah Agr. Expt. Sta. Bull.* 342.
- and —. 1950a. The nutritive value of range forage as affected by vegetation, site, and stage of maturity. *Utah Agr. Expt. Sta. Bull.* 344.
- , L. E. HARRIS AND L. A. STODDART. 1948. Measuring the nutritive content of a foraging sheep's diet under range conditions. *Jour. Animal Sci.* 7(2): 170-180.
- , L. A. STODDART, AND L. E. HARRIS. 1953. Effects of grazing intensity upon the nutritive value of range forage. *Jour. Range Mangt.* 6(1): 51-54.
- DANIEL, H. A. 1934. The calcium, phosphorus, and nitrogen content of grasses and legumes; and the relation of these elements in the plants. *Jour. Amer. Soc. Agron.* 26(6): 496-503.
- AND H. J. HARPER. 1934. The relation between total calcium and phosphorus in mature prairie grass and available plant food in the soil. *Jour. Amer. Soc. Agron.* 26(12): 986-992.
- FERGUSON, W. S. 1931. The seasonal variation in the mineral content of pasture with particular reference to drought. *Jour. Agr. Sci.* 21(2): 233-240.
- GORDON, A. AND A. W. SAMPSON. 1939. Composition of common California foothill plants as a factor in range management. *Calif. Agr. Expt. Sta. Bull.* 627.
- GUILBERT, H. R., S. W. MEAD AND H. C. JACKSON. 1931. The effect of leaching on the nutritive value of forage plants. *Hilgardia* 6(1): 13-25.
- HART, G. H., H. R. GUILBERT AND H. GOSS. 1932. Seasonal changes in the chemical composition of range forage and their relation to nutrition of animals. *Calif. Agr. Expt. Sta. Bull.* 543.
- HOLTZ, H. E. 1930. Effect of calcium and phosphorus content of various soil series in western Washington upon the calcium and phosphorus composition of oats, red clover, and white clover. *Wash. Agr. Expt. Sta. Bull.* 243.
- HULL, A. C. AND J. F. PECHANEC. 1947. Cheatgrass—a challenge to range research. *Jour. Forestry* 45(8): 555-564.
- MAYNARD, L. A. 1937. Interpretations of variations in plant composition in relation to feeding value. *Jour. Amer. Soc. Agron.* 29(3): 504-511.
- . 1947. *Animal nutrition*. McGraw-Hill Book Co., N. Y. 2nd. ed.
- MCCREARY, O. 1927. Wyoming forage plants and their chemical composition. *Studies No. 8. Wyo. Expt. Sta. Bull.* 157: 91-105.
- ORR, J. B. 1929. Minerals in pastures and their relation to animal nutrition. H. K. Lewis & Co., London.
- PATTON, A. R. AND L. GISESEKER. 1942. Seasonal changes in lignin and cellulose content of some Montana grasses. *Jour. Animal Sci.* 1(1): 22-26.
- ROBERTS, E. N. 1926. Wyoming forage plants and their chemical composition. *Studies No. 7. Wyo. Agr. Expt. Sta. Bull.* 146: 35-39.
- SAMPSON, A. W. AND H. E. MALMSTEN. 1926. Grazing periods and forage production on the national forests. *U. S. Dept. Agr. Bull.* 1405.
- SAVAGE, D. A. AND V. G. HELLER. 1947. Nutritional qualities of range forage plants in relation to grazing with beef cattle on the Southern Plains Experimental Range. *U. S. Dept. Agr. Tech. Bull.* 943.
- SCOTT, S. G. 1929. Phosphorus deficiency in forage feeds of range cattle. *Jour. Agr. Res.* 38(2): 113-130.
- SMITH, A. D., R. B. TURNER AND G. A. HARRIS. 1956. The apparent digestibility of lignin by mule deer. *Jour. Range Mangt.* 9(3): 142-145.
- STODDART, L. A. 1941. Chemical composition of *Symphoricarpos rotundifolius* as influenced by soil, site and date of

## NUTRITIVE VALUE OF RANGE FORAGE

225

- collection. Jour. Agr. Res. 63(12): 727-739.
- AND J. E. GREAVES, 1942. The composition of summer range plants in Utah. Utah. Agr. Expt. Sta. Bull. 305.
- AND A. D. SMITH. 1955. Range management. McGraw-Hill Book Co., N. Y. 2d. ed.
- WATKINS, J. M. 1940. The growth habits and chemical composition of brome-grass (*Bromus inermis*), as affected by different environmental conditions.
- Jour. Amer. Soc. Agron. 32(7): 527-538.
- WHITMAN, W. C., *et al.* 1951. Carotene, protein and phosphorus in range and tame grasses of western North Dakota. N. Dak. Agr. Expt. Sta. Bull. 370.