

**SWCA**<sup>®</sup>

ENVIRONMENTAL CONSULTANTS

Sound Science. Creative Solutions.<sup>®</sup>

# **Natural Resources and Wetland Determination Report for the NST Express Pipeline System, McKenzie County, North Dakota**

Prepared for

**NST Express, LLC**

Prepared by

**SWCA Environmental Consultants**

May 2015



**Natural Resources and Wetland Determination Report for the  
NST Express Pipeline System, McKenzie County, North Dakota**

Prepared for:

**NST Express, LLC**

Prepared by:

**Dillon Belisle, B.S., and Kate Kenninger, M.S.  
Environmental Specialists**

Reviewed by:

**Jeff Towner  
Natural Resources Lead**

**SWCA Environmental Consultants  
116 North 4th Street, Suite 200  
Bismarck, North Dakota 58501  
Phone (701) 258-6622, Fax (701) 258-5957**

SWCA Project No. 30320

**May 28, 2015**



**TABLE OF CONTENTS**

	<u>Page</u>
1.0 INTRODUCTION .....	1
1.1 Background .....	1
1.2 Regulatory Background .....	1
1.2.1 Clean Water Act, Section 404 .....	1
1.2.2 USACE Nationwide Permit 12 .....	1
1.2.3 USACE Regional Conditions .....	2
2.0 METHODS .....	2
2.1 Survey area .....	2
2.2 Wetlands .....	4
2.2.1 Hydrophytic Vegetation .....	4
2.2.2 Wetland Hydrology .....	4
2.3 Waterbodies .....	5
2.4 Tree, Sapling, and Shrub Count .....	5
2.5 Noxious Weed Surveys .....	5
2.6 Wildlife Including Threatened and Endangered Species .....	6
2.7 Mapping .....	6
3.0 RESULTS .....	6
3.1 Vegetation .....	6
3.1.1 Herbaceous Upland .....	7
3.1.2 Shrubland and Woody Vegetation .....	7
3.1.3 Cropland .....	7
3.1.4 Hydrophytic Vegetation .....	7
3.2 Hydrology .....	7
3.3 Wetlands .....	8
3.4 Waterbodies .....	9
3.5 Soils .....	9
3.5.1 Williams .....	11
3.5.2 Zahl .....	11
3.5.3 Temvik .....	11
3.5.4 Wilton .....	12
3.5.5 Havrelon .....	12
3.6 Tree, Sapling, and Shrub Count .....	12
3.7 Noxious Weeds .....	13
3.8 Wildlife .....	13
3.8.1 Black-footed Ferret .....	14
3.8.2 Gray Wolf .....	14
3.8.3 Whooping Crane .....	14
3.8.4 Interior Least Tern .....	15
3.8.5 Pallid Sturgeon .....	16
3.8.6 Piping Plover .....	17
3.8.7 Designated Critical Habitat of Piping Plover .....	18
3.8.8 Dakota Skipper .....	18
3.8.9 Rufa Red Knot .....	19
3.8.10 Sprague’s Pipit .....	19

**TABLE OF CONTENTS (continued)**

	<u>Page</u>
3.8.11 Northern Long-eared Bat.....	20
3.8.12 Migratory Birds.....	21
3.8.13 Bald Eagle.....	21
3.8.14 Golden Eagle.....	22
3.8.15 Wildlife Observed.....	22
4.0 CONCLUSIONS AND RECOMMENDATIONS.....	23
5.0 LITERATURE CITED.....	25

**LIST OF TABLES**

<u>Table</u>	<u>Page</u>
1 Monthly Recorded Rainfall at National Weather Service Station in Williston, North Dakota.....	8
2 PEM Wetland Acreage within the Survey Area.....	8
3 Streams within the Survey Area.....	9
4 NRCS Derived Soil Series Present within the Survey Area.....	10
5 Tree, Sapling, and Shrub Count.....	12
6 Wildlife Observed during Field Surveys of the Proposed Pipeline Route.....	22

**LIST OF FIGURES**

<u>Figure</u>	<u>Page</u>
1 Overview of the general topography towards the eastern portion of the pipeline corridor, facing west (photo taken March 20, 2015).....	3
2 Overview of the general topography towards the western portion of the pipeline corridor, facing east (photo taken March 15, 2015).....	3

**LIST OF APPENDICES**

<u>Appendix</u>
A Vicinity Maps and Site Layout Maps
B Survey Area Soil Series Map
C Photographs of Project Corridor

## **1.0 INTRODUCTION**

### **1.1 BACKGROUND**

NST Express, LLC (NST Express) is proposing to construct and operate the proposed NST Express oil pipeline system in McKenzie County, North Dakota. SWCA Environmental Consultants (SWCA) conducted natural resources field surveys in order to identify exclusion and avoidance areas as specified in North Dakota Administrative Code 69-06-08-02 for the proposed NST Express pipeline project.

As proposed, the NST Express pipeline is approximately 23.5 miles long, spanning privately owned lands, in North Dakota (Appendix A). NST Express proposes to bore the Yellowstone River, which is classified as a jurisdictional waterway under Section 10 of the Rivers and Harbors Act (33 United States Code [USC] 403).

SWCA conducted field surveys of a 200-foot-wide survey corridor (100-foot-wide right-of-way [ROW]), including reroutes, a 2-acre survey block of the Fairview rail facility, and a 42-acre block of the proposed Alexander facility on January 30; March 15, 17, 20; and May 1, 2015, to determine the potential presence and extent of wetlands and waterbodies, including jurisdictional waters of the U.S., within the proposed survey area. Concurrently with the wetland determinations, SWCA conducted a cursory threatened and endangered species survey and habitat assessment; a tree, sapling, and shrub enumeration survey; and a noxious weed survey. Site layout maps of the survey area and natural resource features identified during the field surveys are provided in Appendix A.

This report outlines the methodology used by SWCA's biologists to complete each of the aforementioned surveys. Additionally, this report presents the results of the completed field surveys and regulatory recommendations to ensure compliance with the North Dakota Public Service Commission and the U.S. Army Corps of Engineers (USACE) Nationwide Permit 12.

### **1.2 REGULATORY BACKGROUND**

#### **1.2.1 Clean Water Act, Section 404**

Section 404 of the Clean Water Act prohibits the discharge of dredged or fill material into waters of the U.S., also known as jurisdictional waters, without a permit from the USACE.

#### **1.2.2 USACE Nationwide Permit 12**

The USACE Nationwide Permit 12 authorizes the construction of utility lines and associated facilities in waters of the U.S., provided the activity does not result in the permanent loss of greater than 0.5 acre of waters of the U.S., including wetlands.

Nationwide Permit 12 requires that the permittee submit a pre-construction notification prior to commencing construction if any of the following criteria are met.

- The activity involves mechanized land clearing in a forested wetland.
- A Section 10 permit is required to cross a navigable waterbody (Rivers and Harbors Act). Compliance with Section 10 of the Rivers and Harbors Act (33 USC 403) is

required for the passage under the Yellowstone River.

- The utility line exceeds 500 feet in length through any single crossing of a water of the U.S.
- The utility line is placed within a jurisdictional area (i.e., water of the U.S.) and it runs parallel to a stream bed that is within that jurisdictional area.
- Discharges result in the permanent loss of greater than 0.1 acre of waters of the U.S.
- Permanent access roads are constructed above grade in waters of the U.S. for a distance of more than 500 feet.
- Permanent access roads are constructed in waters of the U.S. with impervious materials.

### **1.2.3 USACE Regional Conditions**

The USACE has published several regional conditions for projects operating under Nationwide Permits in North Dakota. The regional conditions apply to wetlands classified as “fens,” waters adjacent to natural springs, the Missouri River, historic properties, and fish spawning areas.

## **2.0 METHODS**

### **2.1 SURVEY AREA**

Overall, northwest North Dakota is characterized by a moderate to cool climate, with cold, dry winters and mild to warm summers. Mean annual precipitation for the area is 14 to 16 inches (Bryce et al. 1998).

The proposed project is located in the Great Plains (level I ecoregion), the West-Central Semi-Arid Prairies (level II ecoregion), and the Northwestern Great Plains (level III ecoregion). Further, the western extent of the alignment is located in the River Breaks (level IV ecoregion) and the eastern extent of the alignment is located in the Missouri Plateau (level IV ecoregion).

The Northwestern Great Plains encompasses the Missouri Plateau section of the Great Plains and consists of semiarid rolling plains of shale, siltstone, and sandstone punctuated by buttes and badlands (Bryce et al. 1998) (Figures 1 and 2). The River Breaks ecoregion forms broken terraces and uplands that descend to the Missouri River and its major tributaries (Bryce et al. 1998). The Missouri Plateau ecoregion was largely unaffected by glaciation and retains its original soils and complex stream drainage pattern (Bryce et al. 1998).



**Figure 1. Overview of the general topography towards the eastern portion of the pipeline corridor, facing west (photo taken March 20, 2015).**



**Figure 2. Overview of the general topography towards the western portion of the pipeline corridor, facing east (photo taken March 15, 2015).**

The inventoried area for the project area discussed herein is situated on the U.S. Geological Survey Fairview (1975), Dore (1971), Buford (1971), Cartwright NE (1971), and Camp Creek West (1979), North Dakota, quadrangles. The proposed NST Express project corridor that was surveyed in 2015 encompasses portions of 25 sections within 4 townships and ranges.

- Section 6, Township (T) 151 North (N), Range (R) 102 West (W)
- Sections 1, 2, 9, 10, 11, 16, 17, 18, and 19, T151N, R103W
- Sections 20, 21, 22, 23, 24, 29, and 30, T151N, R104W
- Sections 23, 24, 26, 31, 32, 33, 34, and 35, T152N, R102W

## **2.2 WETLANDS**

National Wetlands Inventory mapping for the region indicates the presence of wetlands (U.S. Fish and Wildlife Service [USFWS] 2012a). SWCA biologists conducted wetland determinations within the survey area based on the principles and guidelines provided in the *1987 Corps of Engineers Wetlands Delineation Manual* (Manual) (Environmental Laboratory 1987) and the *Regional Supplement to the Corps of Engineers Wetlands Determination Manual: Great Plains Region Version 2.0* (Supplement) (USACE 2010). According to the Manual, an area is a wetland if three mandatory wetland indicators are present in a given area, with special exceptions. These criteria include the presence of hydrophytic vegetation, wetland hydrology, and hydric soils. All wetlands and waterbodies geographically referenced within the survey area during field survey are depicted on the site layout maps in Appendix A.

### **2.2.1 Hydrophytic Vegetation**

Biologists recorded all plants within the vegetative community based on the respective stratum each species occupied. A tree is defined by the Supplement to be a woody-stemmed plant with a trunk diameter at breast height (DBH) of equal to or greater than 3 inches, regardless of height. The sapling and shrub stratum is defined by the Supplement to be composed of woody-stemmed plants with a trunk DBH of less than 3 inches, regardless of height. The herbaceous stratum includes all non-woody-stemmed plants regardless of height. Finally, the woody vine stratum includes all woody-stemmed vines, regardless of diameter.

SWCA recorded the binomial scientific name and percent cover of all plants within a 30-foot radius for the tree stratum, a 15-foot radius for the sapling/shrub stratum, a 5-foot radius for the herbaceous stratum, and a 30-foot radius for the woody vine stratum. SWCA biologists noted each plant species' respective USFWS indicator status (i.e., upland [UPL], facultative upland [FACU], facultative [FAC], facultative wetland [FACW], and obligate [OBL]). Vegetation communities met the hydrophytic vegetation criterion for wetlands if greater than 50% of dominant species had an indicator status of FAC, FACW, and OBL.

### **2.2.2 Wetland Hydrology**

A wetland was determined to contain wetland hydrology if at least one primary indicator or at least two secondary indicators of wetland hydrology were present, as defined by the Manual and Supplement. Common hydrologic indicators include the presence of surface water, high

water table, soil saturation, water marks on trees or other objects, sediment deposits, water-stained leaves, and oxidized rhizospheres on living roots.

### **2.3 WATERBODIES**

Waterbodies (i.e., ponds, creeks, streams, lakes) were identified by the presence of an ordinary high water mark (OHWM). Common identifiable indicators of an OHWM include open water or evidence of a clear, natural line visible on the bank; shelving; changes in soil characteristics; the destruction of terrestrial vegetation; the presence of litter and debris; and watermarks on structures that are inundated during normal high water conditions. The OHWM typically represents the potential limits of the USACE jurisdiction. Please note that the USACE has full discretion in determining the jurisdictional status of referenced wetlands and waterbodies.

SWCA classified streams as perennial, intermittent, or ephemeral based on field observations. During a typical year, a perennial stream contains flowing water year-round and the water table is located above the stream bed. Groundwater is the primary water source for stream flow while precipitation runoff is supplemental. Biologists classified streams that showed significant flow during the field survey as perennial. Additionally, the U.S. Geological Survey topographic maps were used as reference.

An intermittent stream has flowing water for only portions of the year, when groundwater provides water for stream flow. During dry periods, intermittent streams may not have flowing water. Runoff from rainfall is a supplemental source of water for stream flow.

An ephemeral stream has flowing water only during, and for a short duration after, precipitation events in a typical year. Ephemeral stream beds are located above the water table year-round. Groundwater is not a source of water for the stream. Runoff from rainfall is the primary source of water for stream flow.

### **2.4 TREE, SAPLING, AND SHRUB COUNT**

SWCA biologists determined the total number of trees, saplings, and shrubs present within the survey area using several different techniques depending on the type of woody vegetation habitat (i.e., forested upland, shrubland, or shelterbelt) encountered and the overall extent of each habitat within the ROW. The boundary of all forested upland, shrubland, and shelterbelt habitat was geographically referenced using a Trimble GeoXT series handheld global positioning system (GPS) unit. In forested upland and shrubland habitat, SWCA counted the number of all woody-stemmed vegetation with a DBH of  $\geq 1$  inch. In shelterbelt areas, all woody-stemmed vegetation, regardless of DBH, was inventoried via direct count. Biologists taxonomically identified all recorded individuals to the species level within each habitat type.

### **2.5 NOXIOUS WEED SURVEYS**

SWCA conducted a noxious weed survey of all populations of North Dakota state- or county-listed noxious weeds within the project area. Surveys were conducted early in the growing season and it is possible that small isolated patches of noxious weeds could have been missed

or could emerge in other areas in the spring. NST Express will monitor and control noxious weeds within their ROW prior to and subsequent to construction.

## **2.6 WILDLIFE INCLUDING THREATENED AND ENDANGERED SPECIES**

Prior to conducting field surveys, SWCA reviewed information obtained from the USFWS list of threatened and endangered species by North Dakota county (USFWS 2014a) regarding the presence of threatened or endangered species that may occur within the survey area. This document does not represent a comprehensive survey, but rather acknowledges the past and/or current presence of listed species. The lack of discovery of threatened or endangered species does not signify their non-existence within the area, but only that no primary or secondary indications of these species were recorded. SWCA completed a random survey for all listed species and suitable habitat.

A line-of-sight binocular survey for raptor species was also conducted for a distance of approximately 0.5 mile. SWCA biologists noted all wildlife observed during the field survey. Wildlife sightings can involve primary observations (i.e., actual sighting of an animal) or secondary observations (i.e., observation of scat, tracks, or fur deposits).

## **2.7 MAPPING**

The boundaries of each wetland, waterbody, woody vegetation habitat, and noxious weed assemblage were geographically recorded using a Trimble GeoXT GPS unit. The aforementioned GPS unit is capable of recording geographic data with sub-meter accuracy. SWCA used Universal Transverse Mercator Zone 13 North as the projected coordinate system and North American Datum 1983 as the datum. ArcGIS v10.0 (ESRI Redlands, California) was used to analyze recorded features, calculate areas, and generate the maps provided in Appendix A and B. Please note that all data collected using the GPS unit, and displayed on the attached maps, are for review purposes only and do not represent a professional civil survey.

# **3.0 RESULTS**

## **3.1 VEGETATION**

During the field survey, SWCA biologists identified four general types of vegetative communities within the survey area. These vegetative communities were classified as herbaceous upland, shrubland and upland woody vegetation, cropland, and palustrine emergent (PEM) wetland. PEM wetlands are characterized by the presence of herbaceous hydrophytic or submergent aquatic macrophytes. Photographs of the survey area are provided in Appendix C.

Vegetation communities met the hydrophytic vegetation criterion for wetlands if greater than 50% of dominant species had an indicator status of FAC, FACW, or OBL. The upland communities failed to meet at least one of the three assessed wetland criteria.

### 3.1.1 Herbaceous Upland

The herbaceous upland community consists of areas dominated by non-woody vegetation such as grasses and forbs. Two types of herbaceous upland communities were observed during the survey. The dominant herbaceous community within the proposed pipeline survey area includes non-native grasses and forbs including smooth brome (*Bromus inermis*), crested wheatgrass (*Agropyron cristatum*), and Kentucky bluegrass (*Poa pratensis*). The remaining herbaceous upland consisted of mixed-grass prairie comprised of blue grama (*Bouteloua gracilis*), little bluestem (*Schizachyrium scoparium*), junegrass (*Koeleria macrantha*), purple coneflower (*Echinacea angustifolia*), and common yarrow (*Achillea millefolium*).

### 3.1.2 Shrubland and Woody Vegetation

Shrubland communities occurring throughout the survey area consisted of upland areas dominated by woody-stemmed vegetation including primarily silver buffaloberry (*Shepherdia argentea*).

Forested upland vegetation consisted of eastern cottonwood (*Populus deltoides*), Siberian elm (*Ulmus pumila*), green ash (*Fraxinus pennsylvanica*), and Russian olive (*Elaeagnus angustifolia*).

### 3.1.3 Cropland

Cropland was confirmed in the survey area and classified as wheat (*Triticum aestivum*). Other common crops in the area include sugar beets (*Beta vulgaris*), canola (*Brassica napus*), and soybeans (*Glycine max*).

### 3.1.4 Hydrophytic Vegetation

Aquatic vegetation species confirmed during the survey included prairie cordgrass (*Spartina pectinata*), curly dock (*Rumex occidentalis*), Baltic rush (*Juncus balticus*), American sloughgrass (*Beckmannia syzigachne*), and broad-leaf cattail (*Typha latifolia*).

## 3.2 HYDROLOGY

Wetland communities observed during the determination effort displayed at least one primary or two secondary indicators of wetland hydrology, as defined by the Manual and Supplement. Upland communities either failed to display hydrologic indicators or failed to meet the hydrophytic vegetation and hydric soils criterion, as defined by the Manual and Supplement. In some instances, the presence of above average precipitation obscured the wetland/waterbody boundary and OHWM usually present during normal hydrologic conditions. Common indicators of wetland hydrology observed during field surveys include Surface Water (A1), Inundation Visible on Aerial Imagery (B7), Drainage Patterns (B10), Saturation Visible on Aerial Imagery (C9), and Geomorphic Position (D2)

According to National Weather Service preliminary climatological data for Williston, North Dakota, 2.34 inches of precipitation were recorded from November 1, 2014, through May 1, 2015 (Table 1). This amount is 1.72 inches below normal for this time period. Williston is approximately 12 miles north of the project area.

**Table 1. Monthly Recorded Rainfall at National Weather Service Station in Williston, North Dakota**

Month	Recorded Precipitation (inches)	Normal Precipitation (inches)	Difference (inches)
November 2014	0.63	0.65	-0.02
December 2014	0.03	0.62	-0.59
January 2015	0.48	0.59	-0.11
February 2015	0.46	0.39	0.07
March 2015	0.47	0.71	-0.24
April 2015	0.27	1.00	-0.73
May 1, 2015	0.00	0.10	-0.10
<b>Total</b>	<b>2.34</b>	<b>4.06</b>	<b>-1.72</b>

Source: National Oceanic and Atmospheric Administration 2014.

Although precipitation data are not available for the exact project area, it is likely analogous to the precipitation data for Williston for that same timeframe.

### 3.3 WETLANDS

SWCA recorded seven PEM wetlands within the survey area, totaling approximately 1.50 acres. In total, approximately 0.58 acre of PEM wetland is within the 100-foot-wide construction ROW (Table 2). All recorded wetlands are potentially jurisdictional; however, the USACE has the final authority to determine jurisdictional status.

**Table 2. PEM Wetland Acreage within the Survey Area**

Feature ID	Type	USACE Jurisdiction*	Total Recorded Size (acres)	Wetland Area within 100-foot-wide ROW (acres)	Length of Required Crossing (feet)
WET1	Seasonal	Likely Jurisdictional	0.32	0.00	0
WET2	Semipermanent	Likely Jurisdictional	0.07	0.03	17
WET3	Seasonal	Likely Jurisdictional	0.13	0.05	25
WET4	Semipermanent	Likely Jurisdictional	0.28	0.14	66
WET5	Seasonal	Likely Jurisdictional	0.25	0.12	70
WET6	Seasonal	Likely Jurisdictional	0.14	0.08	43
WET7	Seasonal	Likely Jurisdictional	0.31	0.16	76
<b>Total</b>			<b>1.50</b>	<b>0.58</b>	

\*The USACE has the final authority on the jurisdictional status of a waterbody.

ROW = right-of-way

USACE = U.S. Army Corps of Engineers

### 3.4 WATERBODIES

SWCA identified five waterbodies, all classified as intermittent streams, within the survey area (Table 3). All recorded streams are potentially jurisdictional; however, the USACE has the final authority to determine jurisdictional status. STR1, STR2, STR3, STR4 are associated with the Yellowstone Irrigation Canal System, and NST Express proposes to bore these streams.

Upland data points were recorded at potential high risk run-off areas which did not show an OHWM and contained dominantly upland vegetation (Appendix A). However, these areas should be considered when developing a stormwater pollution prevention plan.

NST Express proposes to bore the pipeline beneath the Yellowstone River, which is classified as a jurisdictional waterway under Section 10 of the Rivers and Harbors Act. Nationwide Permit 12 requires that the permittee submit a pre-construction notification prior to commencing construction for passage under a Section 10 waterway.

**Table 3. Streams within the Survey Area**

<b>Feature ID</b>	<b>Description</b>	<b>USACE Jurisdiction*</b>	<b>Total Area (acres)</b>	<b>100-foot-wide ROW Temporary Impact (acres)</b>	<b>Length of Required Crossing (feet)</b>
STR1	Intermittent	Likely Jurisdictional	0.05	0.03	11
STR2	Intermittent	Likely Jurisdictional	0.06	0.03	15
STR3	Intermittent	Likely Jurisdictional	0.09	0.04	20
STR4	Intermittent	Likely Jurisdictional	0.08	0.04	17
STR5	Intermittent	Likely Jurisdictional	0.11	0.05	25
<b>Total</b>			<b>0.39</b>	<b>0.19</b>	

\* The USACE has the final authority on the jurisdictional status of a waterbody.

ROW = right-of-way

USACE = U.S. Army Corps of Engineers

### 3.5 SOILS

Based on Natural Resources Conservation Service (NRCS) mapping (NRCS 2014), 31 soil types are present in the project construction corridor and proposed facilities. The project area analyzed for soils covers the 100-foot-wide construction corridor and proposed Fairview facility expansion and proposed Alexander facility. Table 4 lists all soil units within the project area. The following soil component descriptions represent the most prevalent soil series found within the survey area (NRCS 2014).

**Table 4. NRCS Derived Soil Series Present within the Survey Area**

Soil Types	Map Unit Symbol	Slopes (%)	Acres within 100-foot-wide ROW	Percent within Map Unit
<b>NST Express Pipeline 100-foot-wide Construction Corridor</b>				
Williams-Zahl loams	E3541C	6 to 9	35.61	13.02
Williams-Zahl loams	E3541B	3 to 6	27.70	10.12
Temvik-Wilton-Williams silt loams	E3763B	3 to 6	23.67	8.65
Havrelon silty clay, slightly wet, occasionally flooded	E4132A	0 to 2	21.14	7.72
Temvik-Wilton silt loams	E3755A	0 to 3	17.19	6.28
Temvik-Williams silt loams	E3761A	0 to 3	14.62	5.34
Zahl-Cabba-Maschetah complex	E3609F	6 to 70	13.41	4.90
Zahl-Williams loams	E3555D	9 to 15	13.36	4.88
Havrelon loam, occasionally flooded	E4121A	0 to 2	12.95	4.73
Havrelon loam, slightly wet, occasionally flooded	E4122A	0 to 2	12.46	4.55
Zahl-Williams-Cabba complex	E3639C	6 to 9	11.55	4.22
Zahl-Max loams, dissected	E3567F	15 to 45	9.23	3.37
Temvik-Zahl complex	E3769B	3 to 6	7.84	2.87
Beisigl-Flasher-Telfer loamy fine sands	E1403D	6 to 15	7.32	2.68
Williams-Bowbells loams	E3527B	3 to 6	7.03	2.57
Cabba-Badland complex	E3107F	6 to 70	6.36	2.33
Zahl-Cabba-Williams complex	E3641D	9 to 15	6.27	2.29
Zahl-Tally-Williams complex	E3635C	6 to 9	5.65	2.07
Farnuf loam	E2120B	2 to 6	5.17	1.89
Williams-Bowbells loams	E3527A	0 to 3	2.43	0.89
Water	E4999	0	2.24	0.82
Lohler silty clay, slightly wet, occasionally flooded	E4106A	0 to 2	2.23	0.81
Ridgelawn silt loam, occasionally flooded	E4153A	0 to 2	1.97	0.72
Arnegard loam	E2107A	0 to 2	1.41	0.51
Zahl-Cabba-Arikara complex	E3607F	9 to 70	1.23	0.45
Zahl-Max loams	E3559E	15 to 25	0.91	0.33
Dooley-Zahl complex	E3703B	3 to 6	0.75	0.27
Belfield-Grail clay loams	E0605A	0 to 2	0.63	0.23
Banks fine sandy loam, slightly wet, occasionally flooded	E4202A	0 to 2	0.56	0.20
Lehr-Bowdle loams	E4542B	2 to 6	0.46	0.17
Zahl-Beisigl-Tally complex	E3637D	9 to 15	0.32	0.12
<b>Total</b>			<b>273.67</b>	<b>100.00</b>

Soil Types	Map Unit Symbol	Slopes (%)	Acres within 100-foot-wide ROW	Percent within Map Unit
<b>Alexander Facility</b>				
Williams-Zahl loams	E3541C	6 to 9	23.88	57.18
Williams-Zahl loams	E3541B	3 to 6	8.59	20.58
Williams-Bowbells loams	E3527B	3 to 6	6.66	15.95
Williams-Bowbells loams	E3527A	0 to 3	2.63	6.29
<b>Total</b>			41.76	100.00
<b>Fairview Facility Expansion</b>				
Havrelon loam, occasionally flooded	E4121A	0 to 2	1.15	70.12
Havrelon silty clay, slightly wet, occasionally flooded	E4132A	0 to 2	0.49	29.88
<b>Total</b>			<b>1.64</b>	<b>100.00</b>

Source: Natural Resources Conservation Service 2014.

ROW = right-of-way

### 3.5.1 Williams

The Williams series consists of very deep, slowly permeable, well-drained soils found on glacial till plains and moraines with slopes at approximately 0% to 35%. The mean annual precipitation found throughout the spatial extent of this soil type is approximately 14 inches and mean annual air temperature is approximately 42 degrees Fahrenheit (°F). This soil type is largely used for cultivation. Native vegetation species common to this soil type include western wheatgrass (*Pascopyrum smithii*), needle and thread (*Hesperostipa comata*), blue grama, and green needlegrass (*Nasella viridula*) (NRCS 2014).

### 3.5.2 Zahl

The Zahl series consists of very deep, slowly permeable, well-drained soils found on glacial till plains, moraines, and valley side slopes at approximately 1% to 60%. The mean annual precipitation found throughout the spatial extent of this soil type is approximately 14 inches and mean annual air temperature is approximately 40°F. This soil type is largely used for rangeland foraging. Native vegetation species common to this soil type include western wheatgrass, little bluestem, and needle and thread (NRCS 2014).

### 3.5.3 Temvik

The Temvik series consists of very deep, moderately permeable, well-drained soils that formed in a silty loess mantle overlying glacial till. These soils are on upland plains with slopes that range from 0% to 15%. The mean annual precipitation found throughout the spatial extent of this soil type is 16 inches and mean annual air temperature is 42°F. Soils are commonly cropped to flax (*Linum usitatissimum*), small grains, and corn (*Zea mays*); some areas are used for hay and pasture. Native vegetation is green needlegrass, needle and thread, western wheatgrass, blue grama, upland sedges (*Carex* sp.), and forbs (NRCS 2014).

### 3.5.4 Wilton

The Wilton series consists of very deep, well-drained soils that formed in a silty loess mantle overlying till. These soils are on uplands and have slopes of 0% to 9%. The mean annual precipitation found throughout the spatial extent of this soil type is 16 inches and the mean annual air temperature is 40°F. Cultivated areas are used for growing small grains including flax and corn. Native grasses include western wheatgrass, green needlegrass, bearded wheatgrass (*Elymus caninus*), junegrass, needle and thread, and a variety of forbs (NRCS 2014).

### 3.5.5 Havrelon

The Havrelon series consists of very deep, well- and moderately well-drained, moderately permeable soils that formed in loamy alluvium. These soils are on floodplains of major streams and tributaries and have slopes of 0% to 6%. The mean annual precipitation found throughout the spatial extent of this soil type is 16 inches and the mean annual air temperature is 42°F. Cultivated areas are used for pasture and for growing small grains, hay, and corn. Some areas are irrigated and cropped to sugar beets, potatoes (*Solanum tuberosum*), corn, and alfalfa (*Medicago sativa*). Native grasses include big bluestem (*Andropogon gerardii*), green needlegrass, and western wheatgrass. Trees, including green ash, eastern cottonwood, boxelder (*Acer negundo*), and chokecherry (*Prunus virginiana*), are along the stream channels (NRCS 2014).

## 3.6 TREE, SAPLING, AND SHRUB COUNT

During SWCA’s field survey, nine tree and shrubland areas were geographically referenced within the survey area. Table 5 summarizes the number of trees SWCA counted that may be impacted by the project as currently proposed. The North Dakota Public Service Commission requires a 2:1 post- to pre-construction mitigation for all trees, saplings, and shrubs impacted during the construction of the proposed pipeline. Therefore, SWCA estimates approximately 44 2-year-old sapling individuals would need to be replanted in order to fulfill the 2:1 mitigation requirement.

**Table 5. Tree, Sapling, and Shrub Count**

Woody Vegetation (WV) ID	Species	Type	Number of Trees		Estimated Mitigation Commitment
			Survey Corridor	100-foot-wide Construction ROW	
WV1	Eastern cottonwood ( <i>Populus deltoids</i> )	Tree	1	0	0
WV2	Russian olive ( <i>Elaeagnus angustifolia</i> )	Tree	40	0	0
WV3	Green ash ( <i>Fraxinus pennsylvanica</i> )	Tree	1	0	0
WV4	Siberian elm ( <i>Ulmus pumila</i> )	Tree	40	6	12

Woody Vegetation (WV) ID	Species	Type	Number of Trees		Estimated Mitigation Commitment
			Survey Corridor	100-foot-wide Construction ROW	
WV5	Silver buffaloberry ( <i>Shepherdia argentea</i> )	Shrub	51	0	0
WV6	Silver buffaloberry	Shrub	15	0	0
WV7	Green ash	Tree	1	1	2
WV8	Silver buffaloberry	Shrub	14	13	26
WV9	Green ash	Tree	2	2	4
<b>Total</b>			<b>165</b>	<b>22</b>	<b>44</b>

ROW = right-of-way

### 3.7 NOXIOUS WEEDS

North Dakota Century Code Chapter 63-01.1 and the North Dakota Department of Agriculture recognize 11 species as noxious weeds. The species include absinth wormwood (*Artemisia absinthium*), Canada thistle (*Cirsium arvense*), diffuse knapweed (*Centaurea diffusa*), leafy spurge (*Euphorbia esula*), musk thistle (*Carduus nutans*), purple loosestrife (*Lythrum salicaria*), Russian knapweed (*Acroptilon repens*), spotted knapweed (*Centaurea stoebe*), yellow toadflax (*Linaria vulgaris*), dalmatian toadflax (*Linaria dalmatica*), and salt cedar (*Tamarix ramosissima*). McKenzie County has five county-listed species including black henbane (*Hyoscyamus niger*), common burdock (*Arctium minus*), houndstongue (*Cynoglossum officinale*), halogeton (*Halogeton glomeratus*), and baby’s breath (*Gypsophila paniculata*). SWCA did not identify any areas of state-listed or county-listed noxious weeds within the survey corridor.

### 3.8 WILDLIFE

Several wildlife species that may exist in McKenzie County are listed as threatened or endangered under the Endangered Species Act (ESA) (16 USC 1531 et seq.). According to the USFWS, ESA-listed endangered species in McKenzie County, North Dakota, include the black-footed ferret (*Mustela nigripes*), gray wolf (*Canis lupus*), whooping crane (*Grus americana*), interior least tern (*Sterna antillarum*), and pallid sturgeon (*Scaphirhynchus albus*). Threatened species include the piping plover (*Charadrius melodus*) and its designated critical habitat, Dakota skipper (*Hesperia dacotae*), and rufa red knot (*Calidris canutus rufa*). In addition, Sprague’s pipit (*Anthus spragueii*) is a candidate species and the northern long-eared bat (*Myotis septentrionalis*) is currently listed as threatened pending final ruling on May 4, 2015 (USFWS 2014a). SWCA conducted a cursory threatened and endangered species survey concurrently with the wetland determination. Biologists did not observe any primary (i.e., actual sighting) or secondary (i.e., tracks, scat, fur) indication of the presence of threatened or endangered species.

### 3.8.1 Black-footed Ferret

**Federal Status:** Endangered

Black-footed ferrets are nocturnal, solitary carnivores of the weasel family that have been largely extirpated from the wild primarily due to range-wide decimation of the prairie dog (*Cynomys* sp.) ecosystem (Kotliar et al. 1999). They have been listed by the USFWS as endangered since 1967, and have been the object of extensive re-introduction programs (USFWS 2013a). Ferrets inhabit extensive prairie dog complexes of the Great Plains, typically composed of several smaller colonies in proximity to one another that provide a sustainable prey base. The *Black-footed Ferret Survey Guidelines for Compliance with the Endangered Species Act* (USFWS 1989) states that ferrets require black-tailed prairie dog (*Cynomys ludovicianus*) towns or complexes greater than 80 acres in size, and towns of this dimension may be important for ferret recovery efforts (USFWS 1988a). Prairie dog towns of this size were not observed during the field survey. Therefore, the black-footed ferret is not expected to be impacted by the proposed project.

### 3.8.2 Gray Wolf

**Federal Status:** Endangered

The gray wolf, listed as endangered in the United States in 1978, was believed extirpated from North Dakota in the 1920s and 1930s, with only sporadic reports from the 1930s to present (Licht and Huffman 1996; USFWS 1978). The presence of wolves in most of North Dakota consists of occasional dispersing animals from Minnesota and Manitoba (Licht and Fritts 1994; Licht and Huffman 1996). Most documented gray wolf sightings within western North Dakota are believed to be young males seeking to establish territory (Hagen et al. 2005). The Turtle Mountain region of north-central North Dakota provides marginal habitat that may be able to support a very small population of wolves. The closest known pack of wolves is the Minnesota population located approximately 17 miles (28 kilometers [km]) from the northeast corner of North Dakota.

The gray wolf uses a variety of habitats that support a large prey base, including montane and low-elevation forests, grasslands, and desert scrub (USFWS 2013b). Due to a lack of forested habitat and distance from Minnesota and Manitoba populations, as well as the troubled relationship between humans and wolves and their vulnerability to being shot in open habitats (Licht and Huffman 1996), the re-establishment of gray wolf populations in North Dakota is unlikely. Additionally, habitat fragmentation may further act as a barrier against wolf recolonization in western North Dakota. The gray wolf is not expected to be impacted by the proposed project.

### 3.8.3 Whooping Crane

**Federal Status:** Endangered

The whooping crane was listed as endangered in 1970 in the United States by the USFWS and in 1978 in Canada. Historically, population declines were caused by shooting and destruction of nesting habitat in the prairies from agricultural development. Current threats to the species include habitat destruction, especially suitable wetland habitats that support breeding and

nesting, as well as feeding and roosting during their fall and spring migration (Canadian Wildlife Service and USFWS 2007).

The July 2010 total wild population was estimated at 383 (USFWS 2013c). There is only one self-sustaining wild population, the Aransas-Wood Buffalo National Park population, which nests in Wood Buffalo National Park and adjacent areas in Canada, where approximately 83% of the wild nesting sites occur (Canadian Wildlife Service and USFWS 2007; USFWS 2013c). McKenzie County, including the project area, is within the primary migratory flyway of whooping cranes.

Whooping cranes probe the soil subsurface with their bills for foods on the soil or vegetation substrate (Canadian Wildlife Service and USFWS 2007). Whooping cranes are omnivores and foods typically include agricultural grains, as well as insects, frogs, rodents, small birds, minnows, berries, and plant tubers. The largest amount of time during migration is spent feeding in harvested grain fields (Canadian Wildlife Service and USFWS 2007). Studies indicate that whooping cranes use a variety of habitats during migration, in addition to cultivated croplands, and generally roost in small palustrine (marshy) wetlands within 0.6 mile (1 km) of suitable feeding areas (Howe 1987, 1989). Whooping cranes have been recorded in riverine habitats during their migration, with eight sightings along the Missouri River in North Dakota (Canadian Wildlife Service and USFWS 2007:18). In these cases, they roost on submerged sandbars in wide, unobstructed channels that are isolated from human disturbance (Armbruster 1990).

It is well-documented that migrating whooping cranes use habitats in the vicinity of the proposed project for roosting and feeding. The project area is located within the migratory corridor for the whooping crane, with the nearest sighting being 432 feet from the central portion of the pipeline corridor east of Fairview (USFWS, M. Tarcha, unpublished data). Suitable whooping crane stop over foraging habitat (i.e., cultivated cropland and wetlands >0.04 hectare) was observed within the survey area. Whooping cranes also use sandbars in large streams such as the Yellowstone and Missouri Rivers as stopover roosting habitat. To mitigate any behavioral disturbance that may occur to migrating whooping cranes, if a whooping crane is sighted within 1 mile of the construction area work should be suspended until the species has vacated the area, typically not more than 2 to 3 days. The USFWS should be contacted if any whooping cranes are observed within one mile of the construction right-of-way. If the proposed activity follows these recommendations, the whooping crane is not expected to be impacted by the proposed project.

#### **3.8.4 Interior Least Tern**

**Federal Status:** Endangered

The interior population of the least tern is listed as endangered by the USFWS (1985a). This bird is the smallest member of the gull and tern family, measuring approximately 9 inches in length. Terns remain near flowing water, where they feed by hovering over and diving into standing or flowing water to catch small fish (USFWS 2013d).

The interior population of least terns breeds in isolated areas along the Missouri, Mississippi, Ohio, Red, and Rio Grande river systems, where they nest in small colonies. From late April

to August, terns nest in a shallow hole scraped in an open sandy area, gravel patch, or exposed flat and bare sandbars along rivers, sand and gravel pits, or lake and reservoir shorelines. The adults continue to care for chicks after they hatch. Least terns in North Dakota often will be found sharing sandbars with the piping plover, a threatened species (USFWS 2013d).

Census data indicate over 8,000 least terns in the interior population. In North Dakota, the least tern is found mainly on the Missouri River from Garrison Dam south to Lake Oahe, and on the Missouri and Yellowstone Rivers upstream of Lake Sakakawea (USFWS 1990a, 2013d). Approximately 100 pairs breed in North Dakota (USFWS 2013d). Details of their migration are not known, but their winter range is reported to include the Gulf of Mexico and Caribbean Islands (USFWS 1990a, 2013d).

Loss of suitable breeding and nesting habitat for terns has resulted from dam construction and river channelization on major rivers throughout the Mississippi, Missouri, and Rio Grande river systems. River and reservoir changes have led to reduced sandbar formation and other shoreline habitats for breeding, resulting in population declines. In addition, other human shoreline disturbances affect the species (USFWS 1990a). Critical habitat has not been designated for the species (USFWS 2013d). Current conservation strategies include identification and avoidance of known nesting areas, public education, and limiting or preventing shoreline disturbances near nests and hatched chicks (USFWS 2013d).

Lake Sakakawea is approximately 6.0 miles from the edge of the project area. Endanger interior least terns are known to use the Yellowstone River, which is in the project area, for breeding and feeding. To minimize or reduce potential impacts of the proposed project on the tern, construction should be scheduled to avoid the breeding season to the extent practicable. If the breeding season cannot be avoided, a qualified biological monitor should be deployed on a daily basis for the duration of the construction phase to assess presence/absence of breeding terns. The USACE, in conjunction with the river bore permit, could require monitoring during the tern breeding season. If terns are observed within 0.5 mile of the construction right-of-way at the Yellowstone River, construction should be suspended, and the USFWS should be notified. HDD would be considered as a low-impact crossing technique to reduce or minimize impacts to breeding birds, if present. The interior least tern is not expected to be impacted by the proposed project by following these recommendations or the USACE conditions of approval associated with the Yellowstone River bore permit.

### **3.8.5 Pallid Sturgeon**

**Federal Status:** Endangered

The pallid sturgeon was listed as endangered in 1990 in the United States by the USFWS (1990b). The primary factor leading to the decline of this species is the alteration of habitat through river channelization, creation of impoundments, and alteration of flow regimes (USFWS 1990b). These alterations within the Missouri River have blocked movements to spawning, feeding, and rearing areas; destroyed spawning habitat; altered flow conditions which can delay spawning cues; and reduced food sources by lowering productivity (USFWS 2007). The fundamental elements of pallid sturgeon habitat are defined as the bottom of swift waters of large, turbid, free-flowing rivers with braided channels, dynamic flow patterns, flooding of terrestrial habitats, and extensive microhabitat diversity (USFWS 1990b).

The pallid sturgeon populations occur in the Missouri River below Fort Peck Dam to the headwaters of Lake Sakakawea and the lower Yellowstone River up the confluence of the Tongue River, Montana (USFWS 2007). This population consists of approximately 136 wild adult pallid sturgeon (USFWS 2007). Hatchery-reared sturgeon have also been stocked since 1998. The pallid sturgeon has been found to use the 15.5 miles (25 km) of riverine habitat that would be inundated by Lake Sakakawea at full pool (Bramblett 1996 per USFWS 2007). Larval pallid sturgeons have also been found to drift into Lake Sakakawea. While the majority of pallid sturgeons are found in the headwaters of Lake Sakakawea, the North Dakota Game and Fish Department has caught and released pallid sturgeon in nets set in 80 to 90 feet of water between the New Town and Van Hook areas. Based on this information, pallid sturgeon could be found throughout Lake Sakakawea (personal communication, email from Steve Krentz, Pallid Sturgeon Project Lead, USFWS, to SWCA, September 3, 2010).

Desktop analysis concluded that suitable habitat may be present in the project area. Potential pollution from construction activities, hydrostatic testing, and pipeline operations is a concern for downstream populations of the pallid sturgeon. Activities associated with the proposed project are not anticipated to adversely affect water quality and subsequently the pallid sturgeon. HDD would be considered as a low and/or no impact alternative to traditional wet trench or open-cut methods to cross the Yellowstone River. A USACE Section 10 permit is being completed for the bore of the Yellowstone River. The pallid sturgeon is not expected to be impacted by the proposed project by following these practices or the USACE conditions of approval associated with the Yellowstone River HDD permit.

### **3.8.6 Piping Plover**

#### **Federal Status: Threatened**

The piping plover is a small shorebird which breeds only in three geographic regions of North America: the Atlantic Coast, the Northern Great Plains, and the Great Lakes. Piping plover populations were federally listed as threatened and endangered in 1985, with the Northern Great Plains and Atlantic Coast populations listed as threatened, and the Great Lakes population listed as endangered (USFWS 1985b).

Plovers in the Great Plains make their nests on open, sparsely vegetated sand or gravel beaches adjacent to alkali wetlands, and on beaches, sand bars, and dredged material islands of major river systems (USFWS 2002, 2012b). The shorelines of lakes of the Missouri River constitute significant nesting areas for the bird. Piping plovers nest on the ground, making shallow scrapes in the sand, which they line with small pebbles or rocks (USFWS 1988b). Anthropogenic alterations of the landscape along rivers and lakes where piping plover nest have increased the number and type of predators, subsequently decreasing nest success and chick survival (USFWS 2002, 2012b). The birds fly south by mid to late August to areas along the Texas coast and Mexico (USFWS 2002). The Northern Great Plains population has continued to decline despite federal listing, with population estimates of 1,500 breeding pairs in 1985 reduced to fewer than 1,100 in 1990. Low survival of adult birds has been identified as a factor (Root et al. 1992). Current conservation strategies include identification and preservation of known nesting sites, public education, and limiting or preventing shoreline disturbances near nests and hatched chicks (USFWS 1988b, 2012b).

Desktop analysis concluded that critical habitat for the threatened piping plover is not present in the project area. However, piping plovers are known to use the Yellowstone River for feeding and breeding, which is in the project area. To minimize or reduce potential impacts of the proposed project on the plover, construction should be scheduled to avoid the breeding season to the extent practicable. If the breeding season cannot be avoided, a qualified biological monitor should be deployed on a daily basis for the duration of the construction phase to assess presence/absence of breeding piping plovers. The USACE, in conjunction with the river bore permit, could require monitoring during the piping plover breeding season. If piping plovers are observed within 0.5 mile of the construction right-of-way at the Yellowstone River, construction should be suspended, and the USFWS should be notified. HDD would be considered as a low-impact crossing technique to reduce or minimize impacts to breeding birds, if present. The piping plover is not expected to be impacted by the proposed project by following these recommendations or the USACE conditions of approval associated with the Yellowstone River HDD permit.

### **3.8.7 Designated Critical Habitat of Piping Plover**

The USFWS has designated critical habitat for the Great Lakes and Northern Great Plains populations of piping plover (USFWS 2002). Designated critical habitat for the piping plover includes 183,422 acres and 1,207.5 river miles of habitat including the shoreline of Lake Sakakawea in Mountrail and McKenzie Counties, North Dakota (USFWS 2002).

The proposed project would not modify, alter, disturb, or affect the shoreline of Lake Sakakawea or the Missouri River, but is within the same watershed as Lake Sakakawea and the Missouri River and bores under the Yellowstone River.

### **3.8.8 Dakota Skipper**

**Federal Status:** Threatened

The Dakota skipper is a small butterfly with a 1-inch wingspan. The male wing ranges from a tawny-orange to brown and the female wing is darker brown with tawny-orange spots and faint white spots (USFWS 2011a). The Dakota skipper was found to be warranted for protection under the ESA, was precluded for higher-priority species in 1995, and was the subject of a proposed rule for listing as threatened under the ESA, in addition to a proposed rule for designation of critical habitat in October 2013 (78 *Federal Register* 63625; 78 *Federal Register* 63573). On October 24, 2014, the USFWS determined a threatened species status for the Dakota skipper, and the final rule became effective November 24, 2014 (USFWS 2014b). Of the three North Dakota units of proposed critical habitat, Dakota unit 10 (Eagle Nest Butte) is located in the southwest corner of the Fort Berthold Indian Reservation in McKenzie County. The other two proposed locations are within the U.S. Forest Service lands to the north of Charlson, North Dakota.

The primary cause for the decline of this species includes the loss of high-quality native prairie habitat due to overgrazing, conversion to agriculture, and disruption of natural prairie fire cycles (USFWS 2011a). ‘Type A’ habitat is low wet-mesic prairie with little topographic relief occurring near-shore glacial lake deposits (Royer and Marrone 1992). Three plants species dominate ‘Type A’ habitat and include wood lily (*Lilium philadelphicum*), bluebell bellflower (*Campanula rotundifolia*), and mountain deathcamas (*Zigadenus elegans*)

(McCabe 1981). ‘Type B’ habitat of the Dakota skipper occurs on rolling terrain over gravelly glacial moraine deposits and is dominated by big bluestem, little bluestem, and needlegrasses (*Stipa* spp.), and may include bluebell bellflower and wood lily (USFWS 2014c). Additionally, ‘Type B’ habitat supports extensive stands of purple coneflower, upright prairie coneflower (*Ratibida columnifera*), and common gaillardia (*Gaillardia artistata*) (USFWS 2014c). During the surveys cultivated, non-native grass prairie was dominant in the proposed project area and included species such as smooth brome, crested wheatgrass, and western wheatgrass. Dakota skippers are not known to occur in the project area. Additionally, field surveys concluded that cultivated, non-native grass prairie is dominant in the project area, and therefore not suitable as Dakota skipper habitat. The nearest proposed critical habitat (Unit 12) is located 32.6 miles east-northeast from the proposed project. The Dakota skipper is not expected to be impacted by the proposed project.

### 3.8.9 Rufa Red Knot

#### **Federal Status:** Threatened

The rufa red knot is a medium-sized shorebird approximately 9 to 11 inches in height with breeding plumage consisting of red around the face and a prominent stripe above the eye, breast, and upper belly, and non-breeding plumage a dusky gray and white (Bureau of Indian Affairs 2014). The USFWS published a proposal to list the rufa red knot as threatened under the ESA in the Federal Register in September 2013 (78 *Federal Register* 60023). On December 11, 2014, the USFWS determined a threatened species status for the rufa red knot, and the final rule became effective January 12, 2015 (79 *Federal Register* 73705).

The primary reason for decline of this species includes reduced food supplies in Delaware Bay due to commercial harvest of horseshoe crabs, but also includes areas of range loss due to rising sea levels, shorelines project, and development (USFWS 2013e). The rufa red knot breeds in the Canadian Arctic and migrates 19,000 miles to winter on the U.S. Gulf Coast and in South America. The species generally occurs along the ocean coasts during migration, but a small number have been reported across the interior United States.

Suitable habitat along Lake Sakakawea is approximately 6.0 miles from the edge of the project area, and it is possible the rufa red knot may use the Yellowstone River. However, the likelihood of the rufa red knot occurring in the project area is low. The rufa red knot is not expected to be impacted by the proposed project.

### 3.8.10 Sprague’s Pipit

#### **Federal Status:** Candidate

The Sprague’s pipit is a small passerine, 10 to 15 centimeters in length, endemic to the Northern Great Plains (USFWS 2011b). The Sprague’s pipit requires large tracts of native prairie habitat, unplowed, throughout their life cycle. Because native grasslands are disturbance-dependent, Sprague’s pipit prefers grassland habitats that are regularly disturbed. The frequency of disturbance required for habitat maintenance depends on how quickly grasses grow to an intermediate height (4 to 12 inches) following a disturbance event.

In North Dakota, Sprague's pipit has been found in areas of moderate grazing. Sprague's pipits are sensitive to patch size and avoid edges between grasslands and other habitat features (USFWS 2011b). They may avoid non-grassland features including roads, trails, oil wells, croplands, woody vegetation, and wetlands. The Sprague's pipit is reported to stay up to 350 meters away from anthropogenic features such as roads, oil wells, and wind turbines (USFWS 2011b). The USFWS has estimated that each new oil well and associated road in North Dakota results in potential impacts to approximately 51 acres of pipit habitat due to avoidance and habitat fragmentation (USFWS 2011b). Because of increasing habitat fragmentation, especially by energy development, throughout the Sprague's pipit range, and the loss of native prairie habitat, the Sprague's pipit was listed as a Candidate Species under the ESA in 2010 (USFWS 2011b).

In North Dakota, Sprague's pipit breeds throughout the state except for the easternmost counties. During the breeding season they prefer large patches of well-drained, open native grassland with a minimum size of 358.3 acres (range = 170 to 776 acres). They have not been observed in areas smaller than 71.6 acres on their breeding grounds (USFWS 2011b).

Field surveys concluded that cultivated, non-native grass prairie was dominant in the proposed project area. However, some native grasses of intermediate height occur within the project area. The proposed project is unlikely to directly affect habitat due to lack of adequate patch sizes required by the Sprague's pipit for breeding grounds in the immediate project area, but may indirectly contribute to reduced use of any nearby suitable grassland habitat patches within 350 meters of the proposed project. Sprague's pipit is not expected to be impacted by the proposed project.

### **3.8.11 Northern Long-eared Bat**

#### **Federal Status:** Threatened

On October 2, 2013, the USFWS proposed the northern long-eared bat for listing as endangered under the ESA (USFWS 2013f). This medium-sized bat ranges across the eastern and north central United States and all of the Canadian provinces (USFWS 2013f). Throughout most of this species' range, populations are patchily distributed. They emerge at dusk to fly through the understory of forested hillsides and ridges, feeding on moths, flies, leafhoppers, caddisflies, and beetles.

Most records of northern long-eared bats are from winter hibernacula surveys, with more than 780 hibernacula identified within the United States. No known hibernacula are located in North Dakota, due to either no suitable hibernacula present or a lack of survey effort (USFWS 2013g). This bat species occupies a wide range of rocky and forested habitats. Suitable winter habitat contains large caves and mines (USFWS 2013f). Summer day roosts include abandoned buildings, bridges, hollow trees, stumps, under loose bark, and rock fissures (Jones and Choate 1978). The summer roosting period is from May through October. If woody vegetation is cleared during this period, a qualified biologist should identify any potential roost trees. If potential roost trees are identified further site investigations are recommended.

Suitable winter habitat for northern long-eared bats does not occur in the project area. Nearby trees and rocky outcrops can act as suitable summer day roosts. If these recommendations are followed, the northern long-eared bat is not expected to be impacted by the proposed project.

### 3.8.12 Migratory Birds

**Status:** Protected under the Migratory Bird Treaty Act

Suitable habitat for migratory birds exists in the entire project area. Specifically, grassland nesting birds, waterfowl, wading birds, and shorebirds have the potential to occur, feed, loaf, and nest in the project area, especially during the migratory bird breeding season, which generally occurs between February 1 and July 15. Suitable woodland nesting habitat occurs in the project area, but it is minimal. All take of migratory birds or their active nests, including eggs and young must be avoided to avoid a violation of the Migratory Bird Treaty Act. To minimize or reduce potential impacts of the proposed project on migratory birds, one or a combination of the following should be completed by NST Express.

1. Construction would be conducted to avoid the breeding season (February 1–July 15).
2. Regularly mow and/or clear and grub vegetation in the ROW prior to February 1 and maintain through July 15.
3. Conduct a field survey for nesting migratory birds within 5 days of construction commencement.

If any active nests of migratory birds are discovered, construction should be delayed or halt and the USFWS would be contacted for further direction. If these recommendations are followed, then migratory birds are not expected to be impacted by the proposed project.

### 3.8.13 Bald Eagle

**Federal Status:** Delisted in 2007; protected under the Migratory Bird Treaty Act and the Bald and Golden Eagle Protection Act

The bald eagle (*Haliaeetus leucocephalus*) feeds on fish and carrion and typically roosts in large trees near a water source. Bald eagle nesting habitat is typically any mature stands of conifer (*Pinophyta* sp.) or cottonwood (*Populus* sp.) trees in association with rivers, streams, reservoirs, lakes, or any significant body of water. Bald eagles in North Dakota are usually observed along the Missouri River (Gomes n.d.) and Yellowstone River. Bald eagles frequently migrate through the grassland habitats. One bald eagle was observed during the field surveys; however, no nests were observed during the field surveys. The nearest known eagle nest is approximately 3 miles northwest to the closest portion of the proposed project (North Dakota Game and Fish Department 2015). The USFWS generally recommends a nest buffer of 0.5 mile for any eagles. If any active nests are discovered within 0.5 mile of the proposed pipeline route, construction should halt and the USFWS should be contacted for further direction. If these recommendations are followed, bald eagles are not expected to be impacted by the proposed project.

### 3.8.14 Golden Eagle

**Federal Status:** Unlisted; protected under the Migratory Bird Treaty Act and the Bald and Golden Eagle Protection Act

The golden eagle (*Aquila chrysaetos*) prefers habitat characterized by open prairie, plains, and forested areas. Usually, golden eagles can be found in proximity to badland cliffs which provide suitable nesting habitat. Golden eagles may occur within or near the survey area; however, no golden eagles or nests were observed during the field surveys. The USFWS generally recommends a nest buffer of 0.5 mile for any eagles. If any active nests are discovered within 0.5 mile of the proposed pipeline route, construction should halt and the USFWS should be contacted for further direction. If these recommendations are followed, golden eagles are not expected to be impacted by the proposed project.

### 3.8.15 Wildlife Observed

During the field survey, SWCA biologists observed various wildlife species which use wetlands and other habitat within the survey area (Table 6). Common wildlife species may be affected both directly via death or injury from construction activities or indirectly through the temporary fragmentation of habitat as a result of construction activities and disturbance which may disrupt normal activities such as breeding, feeding, and sheltering.

**Table 6. Wildlife Observed during Field Surveys of the Proposed Pipeline Route**

Common Name	Scientific Name	Observation Type
Rock dove	<i>Columbia livia</i>	Primary
Northern harrier	<i>Circus cyaneus</i>	Primary
Bald eagle	<i>Haliaeetus leucocephalus</i>	Primary
Mule deer	<i>Odocoileus hemionus</i>	Primary
Whitetail deer	<i>Odocoileus virginianus</i>	Primary
Grey partridge	<i>Perdix perdix</i>	Primary
Ringneck pheasant	<i>Phasianus colchicus</i>	Primary
Black-capped chickadee	<i>Poecille atricapillus</i>	Primary
Sharp-tailed grouse	<i>Tympanuchus phasianellus</i>	Primary

## 4.0 CONCLUSIONS AND RECOMMENDATIONS

1. SWCA biologists recorded seven wetlands, totaling approximately 1.50 acres within the 200-foot-wide survey area. No wetlands were observed in the proposed facility areas. In total, approximately 0.58 acre of PEM wetlands is within the 100-foot construction ROW. All may be considered jurisdictional under the USACE standards.
2. SWCA recorded five streams within the survey corridor. All of these exhibited OHWM characteristics and may be considered jurisdictional under the USACE standards.
3. Compliance with Section 10 of the Rivers and Harbors Act (33 USC 403) is required for the passage under the Yellowstone River.
4. SWCA counted 22 trees, saplings, and shrubs that may be impacted by construction activities. Therefore, approximately 44 2-year-old saplings may need to be replanted to fulfill the 2:1 mitigation requirement. According to the recommendations of the North Dakota Forest Service, tree species selection for replacement should be accomplished through collaboration with a reputable area nursery. This will allow for species to be selected based on various factors including species hardiness and area soil type.
5. No threatened or endangered species were observed during the field survey.

Suitable roosting and foraging habitat exists within the project area for the whooping crane. SWCA recommends that if construction is to occur within whooping crane spring and fall migration periods, and a whooping crane is observed within 1 mile of the project, to stop construction and notify the USFWS.

Suitable nesting and foraging habitat for piping plover and interior least tern occurs in the project area along the shorelines of the Yellowstone River. SWCA recommends that construction is conducted outside of the migratory bird nesting season to the extent possible. If construction occurs during the nesting season, the conditions of approval associated with the USACE river bore permit would presumably minimize impacts to the species. Typically, these include monitoring by a qualified biologist or similar BMP.

The Yellowstone River is suitable habitat for pallid sturgeon. NST Express proposes to bore the Yellowstone River.

The other listed threatened and endangered species which occur in McKenzie County are not likely to be detrimentally impacted by construction activities.

6. Migratory birds and habitat were observed throughout the entire project area and a 0.5-mile line-of-sight raptor survey was conducted throughout the survey area.

No active raptor nests were observed, however a northern harrier (*Circus cyaneus*) was observed traversing the survey area. In order to reduce impacts to migratory birds, SWCA recommends conducting construction outside of the migratory bird breeding season as practicable. If construction occurs during the bird breeding season, SWCA recommends to either mow, maintain, or completely remove vegetation within the project construction area outside the migratory bird nesting season, or conduct an avian survey of the project area no greater than 5 days before construction begins. If active nests (i.e., nests with eggs or young) are discovered, the USFWS should be notified.

7. No bald or golden eagle nests were observed. However, one bald eagle individual was observed during the field surveys. If nests (active or not) are discovered within a ½ mile of the proposed activity, the USFWS should be notified.

## 5.0 LITERATURE CITED

- Armbruster, M.J. 1990. Characterization of habitat used by whooping cranes during migration. *Biological Report* 90(4):1–16.
- Bureau of Indian Affairs. 2014. *Programmatic Biological Assessment and Biological Evaluation for Fort Berthold Indian Reservation Oil and Gas Development*. Prepared for the U.S. Fish and Wildlife Service. Prepared by Cardno.
- Bramblett, R.G. 1996. Habitats and movements of pallid and shovelnose sturgeon in the Yellowstone and Missouri Rivers, Montana and North Dakota. Ph.D. dissertation, Montana State University, Bozeman.
- Bryce, S., J.M. Omernik, D.E. Pater, M. Ulmer, J. Schaar, J. Freeouf, R. Johnson, P. Kuck, and S.H. Azevedo. 1998. Ecoregions of North Dakota and South Dakota. Northern Prairie Wildlife Research Center Online. Available at: <http://www.npwrc.usgs.gov/resource/habitat/ndsdeco/index.htm> (Version 30NOV1998). Accessed March 24, 2015.
- Canadian Wildlife Service and U.S. Fish and Wildlife Service (USFWS). 2007. *International Recovery Plan for the Whooping Crane*. Ottawa: Recovery of Nationally Endangered Wildlife (RENEW), and Albuquerque: U.S. Fish and Wildlife Service.
- Environmental Laboratory. 1987. *Corps of Engineers Wetlands Determination Manual*. Technical Report Y-87-1. Vicksburg, Mississippi: U.S. Army Engineer Waterways Experiment Station.
- Gomes, S. n.d. Hawks, eagles, and falcons of North Dakota. North Dakota Game and Fish Department, Bismarck. Northern Prairie Wildlife Research Center Online. Available at: <http://www.npwrc.usgs.gov/resource/birds/hawks/index.htm>. Accessed July 16, 2009.
- Hagen, S.K., P.T. Isakson, and S.R. Dyke. 2005. *North Dakota Comprehensive Wildlife Conservation Strategy*. Bismarck: North Dakota Game and Fish Department.
- Howe, M.A. 1987. Habitat use by migrating whooping cranes in the Aransas-Wood Buffalo corridor. In *Proceedings of the 1985 Crane Workshop*, edited by C. Lewis and J.W. Ziewitz, pp. 303–311. Grand Island, Nebraska: Platte River Whooping Crane Habitat Maintenance Trust and USFWS.
- . 1989. *Migration of Radio-Marked Whooping Cranes from the Aransas-Wood Buffalo Population: Patterns of Habitat Use, Behavior, and Survival*. USFWS Technical Report.
- Jones, J., and J.R. Choate. 1978. Distribution of Two Species of Long-eared Bats of the Genus *Myotis* on the Northern Great Plains. *Prairie Naturalist* 10(2):49–52.


- Kotliar, N.B., B.W. Baker, A.D. Whicker, and G. Plumb. 1999. A Critical Review of Assumptions about the Prairie Dog as a Keystone Species. *Environmental Management* 24(2):177–192.
- Licht, D.S., and S.H. Fritts. 1994. Gray Wolf (*Canis lupus*) Occurrences in the Dakotas. *American Midland Naturalist* 132:74–81.
- Licht, D.S., and L.E. Huffman. 1996. Gray Wolf Status in North Dakota. *The Prairie Naturalist* 28(4):169–174.
- McCabe, T.L. 1981. The Dakota Skipper (*Hesperia dacotae*) (Skinner): Range and Biology, with Special Reference to North Dakota. *Journal of the Lepidopterists' Society* 35(3):179–193.
- National Oceanic and Atmospheric Administration. 2014. Williston and Minot, North Dakota, Preliminary Monthly Climate Data Reports. Available at: <http://www.weather.gov/climate/index.php?wfo=bis>. Accessed May 1, 2014.
- Natural Resources Conservation Service (NRCS). 2014 Web Soil Survey. Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Available at: <http://websoilsurvey.nrcs.usda.gov> and <http://soildatamart.nrcs.usda>. Accessed May 1, 2015.
- North Dakota Game and Fish Department. 2015. North Dakota bald eagle nest sites 2014 data. Received from S. Johnson (NDGF) February 19, 2015.
- Root, B.G., M.R. Ryan, and P.M. Mayer. 1992. Piping Plover Survival in the Great Plains. *Journal of Field Ornithology* 63(1):10–15.
- Royer, R., and G. Marrone. 1992. *Conservation Status of the Dakota Skipper* (*Hesperia dacotae*) in North and South Dakota. U.S. Fish and Wildlife Service, Denver, CO. As cited in 78 *Federal Register* 63573.
- U.S. Army Corps of Engineers (USACE). 2010. *Regional Supplement to the Corps of Engineers Wetland Determination Manual: Great Plains Region Version 2.0*. Edited by J.S. Wakeley, R.W. Lichvar, and C.V. Noble. ERDC/EL TR-08-12. Vicksburg, Mississippi: U.S. Army Engineer Research and Development Center.
- U.S. Fish and Wildlife Service (USFWS). 1978. Reclassification of the gray wolf in the United States and Mexico, with determination of critical habitat in Michigan and Minnesota. *Federal Register* 43(47):9607–9615.
- . 1985a. Interior population of the least tern. *Federal Register* 50 FR 21784–21792. May 28, 1985.
- . 1985b. Endangered and threatened wildlife and plants: determination of endangered and threatened status for the piping plover. *Federal Register* 50(238):50726–50734.

- . 1988a. *Black-footed Ferret Recovery Plan*. Denver: U.S. Fish and Wildlife Service.
- . 1988b. *Great Lakes and Northern Great Plains Piping Plover Recovery Plan*. Twin Cities, Minnesota: U.S. Fish and Wildlife Service.
- . 1989. *Black-footed Ferret Survey Guidelines for Compliance with the Endangered Species Act*. Denver and Albuquerque: U.S. Fish and Wildlife Service.
- . 1990a. *Interior Population of the Least Tern Recovery Plan*. Twin Cities, Minnesota: U.S. Fish and Wildlife Service.
- . 1990b. Endangered and threatened wildlife and plants; Determination of endangered status for the pallid sturgeon. *Federal Register* 55(173):36641–36647.
- . 2002. Designation of critical habitat for the northern Great Plains breeding population of the piping plover; final rule. *Federal Register* 67(176):57637–57717.
- . 2007. *Pallid Sturgeon (Scaphirhynchus albus) 5-year Review Summary and Evaluation*. Billings, Montana: U.S. Fish and Wildlife Service, Pallid Sturgeon Recovery Coordinator.
- . 2011a. Dakota skipper (*Hesperia dacotae*). Bismarck: North Dakota Field Office, Mountain Prairie Region. Available at: [http://www.fws.gov/northdakotafieldoffice/endspecies/species/dakota\\_skipper.htm](http://www.fws.gov/northdakotafieldoffice/endspecies/species/dakota_skipper.htm). Accessed October 22, 2014.
- . 2011b. Endangered and threatened wildlife and plants; 12-month finding on a petition to list Sprague’s pipit as endangered or threatened throughout its range. *Federal Register* 75(178):56028–56050.
- . 2012a. National Wetlands Inventory: Wetlands Online Mapper. Available at: <http://westlandsfew.er.usgs.gov/wtlnds/launch.html>. Accessed July 15, 2012.
- . 2012b. Piping plover. Available at: <http://www.fws.gov/mountain-prairie/species/birds/pipingplover>. Accessed July 15, 2012.
- . 2013a. Black-footed ferret. Available at: <http://ecos.fws.gov/speciesProfile/profile/speciesProfile.action?sPCODE=A004>. Accessed February 21, 2013.
- . 2013b. Gray wolf. Available at: <http://ecos.fws.gov/speciesProfile/profile/speciesProfile.action?sPCODE=A00D>. Accessed February 21, 2013.
- . 2013c. Whooping crane. Available at: <http://ecos.fws.gov/speciesProfile/profile/speciesProfile.action?sPCODE=B003>. Accessed February 21, 2013.

- . 2013d. Least tern (interior population). Available at:  
<http://www.fws.gov/southwest/es/oklahoma/lester.htm>. Accessed February 21, 2013.
- . 2013e. Endangered and Threatened Wildlife and Plants; Proposed Threatened Status for the Rufa Red Knot (*Calidris canutus rufa*). *Federal Register* 78(189):60024–60098.
- . 2013f. *Northern Long-eared Bat Fact Sheet*. Midwest Endangered Species. Available at: <http://www.fws.gov/midwest/endangered/mammals/nlba/nlbaFactSheet.html>. Accessed March 11, 2014.
- . 2013g. Endangered and Threatened Wildlife and Plants; 12-month Finding on a Petition to List the Eastern Small-Footed Bat and the Northern Long-eared Bat as Endangered or Threatened Species; Listing the Northern Long-eared Bat as an Endangered Species; Proposed Rule. *Federal Register* 78(191):61046–61080.
- . 2014a. County occurrence of endangered, threatened, and candidate species and designated critical habitat in North Dakota. Available at:  
[http://www.fws.gov/northdakotafieldoffice/county\\_list.htm](http://www.fws.gov/northdakotafieldoffice/county_list.htm). Accessed April 24, 2014.
- . 2014b. Endangered and Threatened Wildlife and Plants; Threatened Species Status for Dakota Skipper (*Hesperia dacotae*) and Endangered Status for Poweshiek Skipperling (*Oarisma poweshiek*). *Federal Register* 79(206):63672–63748.
- . 2014c. Dakota skipper Section 7 Guidance. U.S. Fish and Wildlife Service, Regions 3 and 6. Available at:  
<http://www.fws.gov/midwest/endangered/insects/dask/pdf/DakotaSkipperS7GuidanceMarch2014.pdf>. Accessed December 1, 2014.

**APPENDIX A**  
**Vicinity Maps and Site Layout Maps**






Service Layer Credits: Sources: Esri, HERE, DeLorme, TomTom, Intermap, iPC, GEBCO, USGS, FAO, NPS, NRCAN, GEBCO, Swisstopo, Esri Japan, METI, Esri China (Hong Kong), Swisstopo, MapmyIndia, © OpenStreetMap contributors, and the GIS User Community

### NST Express Pipeline

- City
- Proposed Pipeline
- U.S. Highway
- State Highway
- Facility Location
- State Boundary
- County Boundary
- Township/Range Boundary


**SWCA**  
 ENVIRONMENTAL CONSULTANTS  
 116 North 4th Street  
 Suite 200  
 Bismarck, ND 58501  
 Phone: 701.258.6622  
 Fax: 701.258.5957  
 www.swca.com



Base Map: 2014 Aerial Imagery  
 Source: USDA/FSA - Aerial Photography Field Office  
 T. 151N, R. 104W & T. 151N, R. 103W &  
 T. 151N, R. 102W & T. 152N, R. 102W  
 McKenzie County, North Dakota



Projection: NAD 1983 UTM Zone 13N




## NST Express Pipeline

- |                       |                   |                         |
|-----------------------|-------------------|-------------------------|
| Upland Data Point     | Existing Road     | NWI Wetland             |
| Proposed Pipeline     | Facility Location | Survey Area             |
| Upland Swale          | Wetland           | Township/Range Boundary |
| Approximate Bore Path | Stream            | Section Boundary        |
| Flowline              | Woody Vegetation  |                         |

**SWCA**  
ENVIRONMENTAL CONSULTANTS


116 North 4th Street  
Suite 200  
Bismarck, ND 58501

Phone: 701.258.6622  
Fax: 701.258.5957  
www.swca.com



Base Map: 2014 Aerial Imagery  
Source: USDA/FSA - Aerial Photography Field Office  
Quadrangle: Fairview (1995), Dore (1991)  
Township/Range: T. 151N, R. 104W  
McKenzie County, North Dakota






## NST Express Pipeline

- |                       |                   |                         |
|-----------------------|-------------------|-------------------------|
| Upland Data Point     | Existing Road     | NWI Wetland             |
| Proposed Pipeline     | Facility Location | Survey Area             |
| Upland Swale          | Wetland           | Township/Range Boundary |
| Approximate Bore Path | Stream            | Section Boundary        |
| Flowline              | Woody Vegetation  |                         |

**SWCA**<sup>®</sup>  
 ENVIRONMENTAL CONSULTANTS


116 North 4th Street  
 Suite 200  
 Bismarck, ND 58501

Phone: 701.258.6622  
 Fax: 701.258.5957  
 www.swca.com



Base Map: 2014 Aerial Imagery  
 Source: USDA/FSA - Aerial Photography Field Office  
 Quadrangle: Dore (1991), Buford (1976)  
 Township/Range: T. 151N, R. 104W  
 McKenzie County, North Dakota






## NST Express Pipeline

- |                       |                   |                         |
|-----------------------|-------------------|-------------------------|
| Upland Data Point     | Existing Road     | NWI Wetland             |
| Proposed Pipeline     | Facility Location | Survey Area             |
| Upland Swale          | Wetland           | Township/Range Boundary |
| Approximate Bore Path | Stream            | Section Boundary        |
| Flowline              | Woody Vegetation  |                         |

**SWCA**  
 ENVIRONMENTAL CONSULTANTS


116 North 4th Street  
 Suite 200  
 Bismarck, ND 58501

Phone: 701.258.6622  
 Fax: 701.258.5957  
 www.swca.com



Base Map: 2014 Aerial Imagery  
 Source: USDA/FSA - Aerial Photography Field Office  
 Quadrangle: Buford (1976)  
 Township/Range: T. 151N, R. 104W & T. 151N, R. 103W  
 McKenzie County, North Dakota






## NST Express Pipeline

- |                       |                   |                         |
|-----------------------|-------------------|-------------------------|
| Upland Data Point     | Existing Road     | NWI Wetland             |
| Proposed Pipeline     | Facility Location | Survey Area             |
| Upland Swale          | Wetland           | Township/Range Boundary |
| Approximate Bore Path | Stream            | Section Boundary        |
| Flowline              | Woody Vegetation  |                         |



116 North 4th Street  
Suite 200  
Bismarck, ND 58501


Phone: 701.258.6622  
Fax: 701.258.5957  
www.swca.com



Base Map: 2014 Aerial Imagery  
Source: USDA/FSA - Aerial Photography Field Office  
Quadrangle: Buford (1976)  
Township/Range: T. 151N, R. 103W  
McKenzie County, North Dakota

Projection: NAD 1983 UTM Zone 13N






## NST Express Pipeline

- |                       |                   |                         |
|-----------------------|-------------------|-------------------------|
| Upland Data Point     | Existing Road     | NWI Wetland             |
| Proposed Pipeline     | Facility Location | Survey Area             |
| Upland Swale          | Wetland           | Township/Range Boundary |
| Approximate Bore Path | Stream            | Section Boundary        |
| Flowline              | Woody Vegetation  |                         |

**SWCA**<sup>®</sup>  
 ENVIRONMENTAL CONSULTANTS


116 North 4th Street  
 Suite 200  
 Bismarck, ND 58501

Phone: 701.258.6622  
 Fax: 701.258.5957  
 www.swca.com



Base Map: 2014 Aerial Imagery  
 Source: USDA/FSA - Aerial Photography Field Office  
 Quadrangle: Buford (1976), Cartwright NE (1991)  
 Township/Range: T. 151N, R. 103W  
 McKenzie County, North Dakota






## NST Express Pipeline

- |                       |                   |                         |
|-----------------------|-------------------|-------------------------|
| Upland Data Point     | Existing Road     | NWI Wetland             |
| Proposed Pipeline     | Facility Location | Survey Area             |
| Upland Swale          | Wetland           | Township/Range Boundary |
| Approximate Bore Path | Stream            | Section Boundary        |
| Flowline              | Woody Vegetation  |                         |

**SWCA**  
ENVIRONMENTAL CONSULTANTS


116 North 4th Street  
Suite 200  
Bismarck, ND 58501

Phone: 701.258.6622  
Fax: 701.258.5957  
www.swca.com



Base Map: 2014 Aerial Imagery  
Source: USDA/FSA - Aerial Photography Field Office  
Quadrangle: Cartwright NE (1991)  
Township/Range: T. 151N, R. 103W  
McKenzie County, North Dakota






## NST Express Pipeline

- |                       |                   |                         |
|-----------------------|-------------------|-------------------------|
| Upland Data Point     | Existing Road     | NWI Wetland             |
| Proposed Pipeline     | Facility Location | Survey Area             |
| Upland Swale          | Wetland           | Township/Range Boundary |
| Approximate Bore Path | Stream            | Section Boundary        |
| Flowline              | Woody Vegetation  |                         |



116 North 4th Street  
Suite 200  
Bismarck, ND 58501

Phone: 701.258.6622  
Fax: 701.258.5957  
www.swca.com



Base Map: 2014 Aerial Imagery  
Source: USDA/FSA - Aerial Photography Field Office  
Quadrangle: Cartwright NE (1991)  
Township/Range: T. 151N, R. 103W & T. 151N, R. 102W  
McKenzie County, North Dakota






## NST Express Pipeline

- |                       |                   |                         |
|-----------------------|-------------------|-------------------------|
| Upland Data Point     | Existing Road     | NWI Wetland             |
| Proposed Pipeline     | Facility Location | Survey Area             |
| Upland Swale          | Wetland           | Township/Range Boundary |
| Approximate Bore Path | Stream            | Section Boundary        |
| Flowline              | Woody Vegetation  |                         |



116 North 4th Street  
Suite 200  
Bismarck, ND 58501

Phone: 701.258.6622  
Fax: 701.258.5957  
www.swca.com



Base Map: 2014 Aerial Imagery  
Source: USDA/FSA - Aerial Photography Field Office  
Quadrangle: Cartwright NE (1991), Camp Creek West (1978)  
Township/Range: T. 151N, R. 102W & T. 152N, R. 102W  
McKenzie County, North Dakota






## NST Express Pipeline

- |                       |                   |                         |
|-----------------------|-------------------|-------------------------|
| Upland Data Point     | Existing Road     | NWI Wetland             |
| Proposed Pipeline     | Facility Location | Survey Area             |
| Upland Swale          | Wetland           | Township/Range Boundary |
| Approximate Bore Path | Stream            | Section Boundary        |
| Flowline              | Woody Vegetation  |                         |




116 North 4th Street  
Suite 200  
Bismarck, ND 58501

Phone: 701.258.6622  
Fax: 701.258.5957  
www.swca.com



Base Map: 2014 Aerial Imagery  
Source: USDA/FSA - Aerial Photography Field Office  
Quadrangle: Camp Creek West (1978)  
Township/Range: T. 152N, R. 102W  
McKenzie County, North Dakota






# NST Express Pipeline

- ▲ Upland Data Point
- Proposed Pipeline
- Upland Swale
- Approximate Bore Path
- - - Flowline
- Existing Road
- Facility Location
- Wetland
- Stream
- Woody Vegetation
- NWI Wetland
- Survey Area
- Township/Range Boundary
- Section Boundary




116 North 4th Street  
Suite 200  
Bismarck, ND 58501  
Phone: 701.258.6622  
Fax: 701.258.5957  
www.swca.com



Base Map: 2014 Aerial Imagery  
Source: USDA/FSA - Aerial Photography Field Office  
Quadrangle: Camp Creek West (1978)  
Township/Range: T. 152N, R. 102W  
McKenzie County, North Dakota






## NST Express Pipeline

- |                       |                   |                         |
|-----------------------|-------------------|-------------------------|
| Upland Data Point     | Existing Road     | NWI Wetland             |
| Proposed Pipeline     | Facility Location | Survey Area             |
| Upland Swale          | Wetland           | Township/Range Boundary |
| Approximate Bore Path | Stream            | Section Boundary        |
| Flowline              | Woody Vegetation  |                         |



116 North 4th Street  
Suite 200  
Bismarck, ND 58501

Phone: 701.258.6622  
Fax: 701.258.5957  
www.swca.com




Base Map: 2014 Aerial Imagery  
Source: USDA/FSA - Aerial Photography Field Office  
Quadrangle: Camp Creek West (1978)  
Township/Range: T. 152N, R. 102W  
McKenzie County, North Dakota



**APPENDIX B**  
**Survey Area Soils Series Map**






# NST Express Pipeline

- Proposed Pipeline
- Soil Unit Boundary
- Section Boundary
- Township/Range Boundary
- Approximate Bore Path
- Survey Area
- Existing Road
- Facility Location




116 North 4th Street  
 Suite 200  
 Bismarck, ND 58501  
 Phone: 701.258.6622  
 Fax: 701.258.5957  
 www.swca.com



Base Map: 2014 Aerial Imagery  
 Source: USDA/FSA - Aerial Photography Field Office  
 Quadrangle: Fairview (1995), Dore (1991)  
 Township/Range: T. 151N, R. 104W  
 McKenzie County, North Dakota






# NST Express Pipeline

- Proposed Pipeline
- Soil Unit Boundary
- Section Boundary
- Approximate Bore Path
- Survey Area
- Township/Range Boundary
- Existing Road
- Facility Location




116 North 4th Street  
Suite 200  
Bismarck, ND 58501  
Phone: 701.258.6622  
Fax: 701.258.5957  
www.swca.com



Base Map: 2014 Aerial Imagery  
Source: USDA/FSA - Aerial Photography Field Office  
Quadrangle: Dore (1991), Buford (1976)  
Township/Range: T. 151N, R. 104W  
McKenzie County, North Dakota








## NST Express Pipeline

- Proposed Pipeline
- Approximate Bore Path
- Existing Road
- Facility Location
- Soil Unit Boundary
- Survey Area
- Township/Range Boundary
- Section Boundary




116 North 4th Street  
Suite 200  
Bismarck, ND 58501  
Phone: 701.258.6622  
Fax: 701.258.5957  
www.swca.com



Base Map: 2014 Aerial Imagery  
Source: USDA/FSA - Aerial Photography Field Office  
Quadrangle: Buford (1976)  
Township/Range: T. 151N, R. 103W  
McKenzie County, North Dakota






## NST Express Pipeline

- Proposed Pipeline
- Approximate Bore Path
- Existing Road
- Facility Location
- Soil Unit Boundary
- Survey Area
- Township/Range Boundary
- Section Boundary




116 North 4th Street  
 Suite 200  
 Bismarck, ND 58501  
 Phone: 701.258.6622  
 Fax: 701.258.5957  
 www.swca.com



Base Map: 2014 Aerial Imagery  
 Source: USDA/FSA - Aerial Photography Field Office  
 Quadrangle: Buford (1976), Cartwright NE (1991)  
 Township/Range: T. 151N, R. 103W  
 McKenzie County, North Dakota






## NST Express Pipeline

- Proposed Pipeline
- Approximate Bore Path
- Existing Road
- Facility Location
- Soil Unit Boundary
- Survey Area
- Township/Range Boundary
- Section Boundary




116 North 4th Street  
Suite 200  
Bismarck, ND 58501  
Phone: 701.258.6622  
Fax: 701.258.5957  
www.swca.com



Base Map: 2014 Aerial Imagery  
Source: USDA/FSA - Aerial Photography Field Office  
Quadrangle: Cartwright NE (1991)  
Township/Range: T. 151N, R. 103W  
McKenzie County, North Dakota






## NST Express Pipeline

- Proposed Pipeline
- Approximate Bore Path
- Existing Road
- Facility Location
- Soil Unit Boundary
- Survey Area
- Township/Range Boundary
- Section Boundary




116 North 4th Street  
Suite 200  
Bismarck, ND 58501  
Phone: 701.258.6622  
Fax: 701.258.5957  
www.swca.com



Base Map: 2014 Aerial Imagery  
Source: USDA/FSA - Aerial Photography Field Office  
Quadrangle: Cartwright NE (1991)  
Township/Range: T. 151N, R. 103W & T. 151N, R. 102W  
McKenzie County, North Dakota






## NST Express Pipeline

- Proposed Pipeline
- Approximate Bore Path
- Existing Road
- Facility Location
- Soil Unit Boundary
- Survey Area
- Township/Range Boundary
- Section Boundary



116 North 4th Street  
Suite 200  
Bismarck, ND 58501  
Phone: 701.258.6622  
Fax: 701.258.5957  
www.swca.com




Base Map: 2014 Aerial Imagery  
Source: USDA/FSA - Aerial Photography Field Office  
Quadrangle: Cartwright NE (1991), Camp Creek West (1978)  
Township/Range: T. 151N, R. 102W & T. 152N, R. 102W  
McKenzie County, North Dakota










## NST Express Pipeline

- Proposed Pipeline
- Approximate Bore Path
- Existing Road
- Facility Location
- Soil Unit Boundary
- Survey Area
- Township/Range Boundary
- Section Boundary



116 North 4th Street  
 Suite 200  
 Bismarck, ND 58501  
 Phone: 701.258.6622  
 Fax: 701.258.5957  
 www.swca.com



Base Map: 2014 Aerial Imagery  
 Source: USDA/FSA - Aerial Photography Field Office  
 Quadrangle: Camp Creek West (1978)  
 Township/Range: T. 152N, R. 102W  
 McKenzie County, North Dakota



**APPENDIX C**  
**Photographs of Project Area**

---





**Figure C.1. Seasonal wetland (WET1), facing southwest (photo taken March 15, 2015).**



**Figure C.2. Single eastern cottonwood (*Populus deltoides*) (WV1), facing west (photo taken March 15, 2015).**



**Figure C.3. Intermittent stream (STR2) associated with irrigation canal, facing south (photo taken March 15, 2015).**



**Figure C.4. Intermittent stream (STR3) associated with irrigation canal, facing south (photo taken March 15, 2015).**



**Figure C.5. Intermittent stream (STR4) associated with irrigation canal, facing north (photo taken March 15, 2015).**



**Figure C.6. Green ash (*Fraxinus pennsylvanica*), facing east (photo taken March 17, 2015).**



**Figure C.7. Seasonal wetland (WET4), facing north (photo taken March 17, 2015).**



**Figure C.8. Semipermanent wetland (WET5), facing north (photo taken March 17, 2015).**



**Figure C.9. Seasonal wetland (WET8), facing north (photo taken March 20, 2015).**



**Figure C.10. Seasonal fringe wetland (WET7), facing north (photo taken March 20, 2015).**



**Figure C.11. Overview of proposed Fairview rail facility expansion area, facing southeast (photo taken May 1, 2015).**



**Figure C.12. Overview of proposed Alexander facility area, facing east (photo taken May 1, 2015).**