

To:
North Dakota Public Service Commission

Commissioners,

Thank you for inviting us today, and thank you for listening. These are items that have generational effects, long after the trainloads of Federal Carbon Credits have been put in the bank, and therefore warrant the attention.

If the SCS pipeline project were for the purpose of distributing power, comm, water, or other benefitting utilities, there would be an entirely different response to this project. However, this proposed hazardous pipeline project is a for-private-profit project, at the expense of the landowners.

Presenter information:

- Mitchell B. Kertzman
- Grew up in Hazelton (less than 2 miles from the proposed route)
- Military Veteran (USAF) 8/30/1977 to 8/31/1996
- Retired Civil Engineer (SDSMT 2001 - B.S. Civil Engineering)
- Represented properties:
 - Township 135 North, Range 76 West, Section 26: E2E2, 5th PM ND (Emmons)
 - Township 135 North, Range 76 West, Section 26: W2E2, 5th PM ND (Emmons)

1. Concerning public safety and well-being; what happens during a CO₂ 'spill' is not the same as what happens during an oil spill:

Carbon dioxide is not the same animal as petrochemicals:

- CO₂ is a natural trace element in our atmosphere.
- CO₂ is not poisonous.
- CO₂ is non-flammable (no fire, no explosion).
- CO₂ is colorless and odorless.
- CO₂ is heavier than the normal mix of atmosphere.



- CO₂ has to be piped at a minimum pressure of 1,100 psi under perfect conditions; nearly double that in reality. Gas pipelines run at around ¼ of that. For comparison, the normal water pressure to your house is 60 psi.

In the event of a pipe break or inadvertent release, the semi-fluid carbon dioxide, upon exiting the confines of the pipe, increases in velocity and decreases in pressure, and phases to gas. There is no explosion or smell (unless an odorant has been added). The CO₂ gas, being heavier than the surrounding normal mix of atmosphere, finds a low spot and fills it to a volume equaling the gas volume resulting from the escaped liquid CO₂. The CO₂ gas displaces the normal atmosphere. this can occur in a flat field or a draw or a lake.

Walking around in a cloud of CO₂ is in itself not a problem for the first few seconds. CO₂ toxicity sets in quickly and cognition ends. The next problem lies in the fact that there is *only* CO₂ to breathe. All creatures with lungs die quickly without oxygen. Indeed, any functioning rescue vehicles would have to be electric powered, as internal combustion engines don't function without oxygen.

By the time local fire/rescue realize there is a CO₂ problem, and muster *non-existent* electric-powered rescue equipment, it'll be too late to save anyone within the bubble of carbon dioxide. Then it's just a matter of waiting for the cloud to disburse into the atmosphere and then go in to gather the bodies.

The survival time in an anoxic environment is approximately equal to the time an average swimmer can stay alive under water. It only takes a 10% concentration of CO₂ to cause convulsions, coma, and death (*Ref: attached IJEM report*). A CO₂ bubble from a failed pipe weld or breach would certainly be around 90% - 95% CO₂.

The situation is compounded by the fact that most won't even know there is a problem, and while they are still conscious, won't know what is happening, what is wrong, or which direction is towards safety.

The cloud of CO₂ eventually disburses back into the atmosphere. The extent of the cloud could be mapped later by surveying the carcass locations.

The group that seeks the trainload of carbon credit money will tell you that the installation and operation of the pipeline is governed by approved safety standards, and that a breach is highly unlikely.

Consider this: SCSs proposed project consists of approximately 1900 miles of pipe. Assuming a laying length of 40', that makes something near 250,800 butt-welds, plus fittings, constructed by the lowest bidders, on a highly accelerated schedule. When a parallel pipeline is added to the route, the likelihood of disaster doubles.

2. Concerning the immediate impact to our farmland:

Because this proposed pipeline route runs diagonally on the map, rights-of-way are ignored, causing excavations that run through the heart of the most valuable farmland. As a project that claims to be good for Agriculture, it does a good job of ruining farmground. SCS claims to provide three years of crop loss relief. That's pretty generous. It will take several generations to restore the topsoil back to a pre Carbon Credit state.

Land value will immediately decrease due to the lowered productivity potential and threat of rupture. The yearly land tax bill will not be reduced. How many people do you know that have said they'd love to buy some land with a CO₂ time bomb on it?

This immediate impact is the least of the problems created.

3. Concerning the generational impact to our land:

For the next 99 years (five generations):

The land, with a pipeline containing a very large amount of asphyxiating gas, will never again be a desirable location to build a home. It will never again be a desirable location for any use other than non-animal agriculture. This fixes the highest use of the land. It becomes a 320-acre tax liability.

Five generations of my family will have to live with the possibility of a catastrophic event that ends the life of people, livestock and wildlife in an area defined, at the time, by wind speed/direction, topology, and volume of CO₂ released.

Five generations of my family will pay property taxes on acres that will never have value greater than cropland. Five generations of my family will pay property taxes on land that has been taken for use by a group of private profiteers, for their monetary gain.

Five generations of my family will have to somehow be prepared to be sued by a harmed party if/when there is a break on our property, as there is no insurance coverage for a CO₂ pipeline break.

The sixth generation will be left with the responsibility of cleaning up the mess when the pipeline is abandoned.

4. Concerning cover depth:

Given the size and weight of modern farm equipment, envision what happens when a massive piece of equipment gets mired in, in a low spot, with a 24" or larger welded steel pipe just 48" below the surface. A perfectly bedded and perfectly welded pipe *might* survive. Damage to the pipe and the consequences of the release of a large volume of CO₂ would be all on the landowner, not the Profiteer.

A cover depth of 48" is inadequate under Ag ground. 6' of cover would be more appropriate. This also gets the majority of the pipe below frost depth, thus somewhat mitigating movement of the high-pressure welded steel pipe. Frost heave generates impressive forces.

5. Closing:

In earlier times, when someone tried to take another's land or livestock for their own benefit, this was considered a hostile act, and people usually died, most often by gunfire, sometimes by hanging, and with the law on the Owners side.

Currently, as it appears, the one wanting to do the taking simply needs investors with very deep pockets, regardless of the Country the investors hail from.

We understand that Summit will likely assume the load from another proposed CO₂ pipeline that has dropped out (Navigator). This will more or less double the currently proposed flow rate, meaning another pipe running parallel to the proposed 24", most likely another 24". Or a single 36". Either way, this would be the largest high-pressure CO₂ pipeline on the planet.

A cover depth of 48" is inadequate under Ag ground. 6' of cover would be more appropriate to mitigate damage from farm equipment. We ask that the Commission consider a stipulation to provide 6' of cover where the pipeline is under Ag land.

If the Commission approves this route, what is to stop SCS from taking whatever they want using Eminent Domain?

Until emergency response training and equipment is clearly defined and bonded, it would be incorrect to approve the route. Training and equipment provided by the Pipeline Operator, with provisions to keep the training and equipment viable and funded for 99 years. This is vital, as the likelihood of pipeline failure increases over time. Can we expect the counties along the route to fund and maintain a CO₂ response team and the required equipment for the next 99 years? For a pipeline that has no benefit to the counties?

The Operator of a potentially deadly private pipeline facility should be responsible for the cost of providing emergency response and disaster cleanup for the life of the facility, not the counties they operate in. Alternately, Consider the utility of a CO₂ pipeline safety tax, paid by the pipeline Owner, to the affected counties, for the purpose of funding training and equipment, for as long as the pipeline is in operation.

Kertzman Farm Trust -and- Dwight Edwin Kertzman and Wanda Lee Kertzman Revocable Living Trust



Mitchell B. Kertzman

Representing these properties:

Township 135 North, Range 76 West, Section 26: E2E2, Emmons County, North Dakota
Kertzman Farm Trust

Mitchell B. Kertzman, Trustee

-and-

Township 135 North, Range 76 West, Section 26: W2E2, Emmons County, North Dakota
Dwight Edwin Kertzman and Wanda Lee Kertzman Revocable Living Trust

As a library, NLM provides access to scientific literature. Inclusion in an NLM database does not imply endorsement of, or agreement with, the contents by NLM or the National Institutes of Health.

Learn more: [PMC Disclaimer](#) | [PMC Copyright Notice](#)

International Journal
of Emergency Medicine

BMC

Int J Emerg Med. 2017; 10: 14.

PMCID: PMC5380556

Published online 2017 Apr 4. doi: [10.1186/s12245-017-0142-y](https://doi.org/10.1186/s12245-017-0142-y)

PMID: [28378268](https://pubmed.ncbi.nlm.nih.gov/28378268/)

Carbon dioxide poisoning: a literature review of an often forgotten cause of intoxication in the emergency department

[Kris Permentier](#),¹ [Steven Vercammen](#),^{✉2} [Sylvia Soetaert](#),² and [Christian Schellemans](#)¹

Abstract

The goal of this article was to provide an overview of the literature available on carbon dioxide intoxication. Articles were included based on their focus on medical or physiological effects of carbon dioxide. Studies related to decompression sickness were excluded. Mechanisms of carbon dioxide poisoning (both as an asphyxiant and as a toxicant) were described. Our review suggested that precautions are needed when handling dry ice or while working in confined spaces. Pre-hospital responders also need to pay attention for the possible diagnosis of CO₂ intoxication for their own safety. When confronted with a victim, he/she should be removed from the dangerous area as fast as possible and oxygen should be administered. Without adequate treatment, victims may show acute reduced cognitive performance, respiratory failure, and circulatory arrest. Therefore, carbon dioxide poisoning is a rare but not to miss diagnosis in the emergency department.

Keywords: Carbon dioxide, Carbon dioxide intoxication, Dry ice, Intoxication

Review

Background

Carbon dioxide (CO₂) is a product of combustion, fermentation, and respiration. In normal air, carbon dioxide percentages are very low (around 0.04%). It is a colorless, odorless, and flammable gas that accumulates near the ground (CO₂ is 1.5 times heavier than air). These characteristics explain why enclosed environments are vulnerable for CO₂ buildup, displacing oxy

[Back to Top](#)

from the area [1]. The term “confined space hypoxic syndrome” has been proposed to describe confined space accidents occurring in water meter pits, tanks, holds of ships, mines, underground storage bins, and so forth, resulting from oxygen-deficient atmospheres [2, 3].

Studies conducted in the 1980s showed that there were 89 deaths per year in the USA alone, during work in confined spaces; 22% occur during rescue efforts [4]. A number has not decreased since. A recent study of the Occupational Safety and Health Administration (OSHA) in 2015 still estimated the number of deaths involving confined spaces to be around 90 per year. Unfortunately, the number of killed rescuers has risen to two thirds of those deaths [5].

Carbon dioxide does not only cause asphyxiation by hypoxia but also acts as a toxicant. At high concentrations, it has been showed to cause unconsciousness almost instantaneously and respiratory arrest within 1 min [6].

Other causes of carbon dioxide intoxication have been identified as well, such as dry ice. Dry ice undergoes sublimation (direct conversion from a solid state into a gas), and if it is warmed rapidly, large amounts of CO₂ are generated, which is especially dangerous in closed environments [7–9]. Dry ice intoxication might be accidental [1, 7, 10, 11] or deliberately as several suicide cases have been described [8, 9]. Massive geothermal emissions have also been described as a possible cause of CO₂ intoxications, though in these cases, a relation with other toxic gasses inhaled might not be excluded [12, 13]. We performed this literature review to understand the contribution of CO₂ specifically to “confined space hypoxic syndrome,” and wanted to make physicians more aware of the condition.

Methods

A literature review was performed where articles were sought in MEDLINE medical database via PubMed using the search terms: “dry ice poisoning,” “carbon dioxide poisoning,” “CO₂ poisoning,” “carbon dioxide intoxication,” and “CO₂ intoxication.” The references of the articles that were found were further evaluated to incorporate as much of the available literature as possible. Articles were included based on their focus on medical or physiological effects of carbon dioxide. Studies related to decompression sickness were excluded. Each article was independently evaluated by two of the authors. For all of the articles, an unanimous consensus among the authors was reached whether to include or exclude. In the end, a total of 19 articles were evaluated to be relevant to this topic and included in this review.

Results

CO₂ toxicity in animal models Tests performed on mongrel dogs show the physiological effect of carbon dioxide on the body: after inhalation of a 50% CO₂ and 50% air mixture, respiratory movement increased for about 2 min, and then, it decreased for 30–90 min. Hill and Flack showed that CO₂ concentrations up to 35% have an exciting effect upon both circulation and respiration, but those beyond 35% are depressant upon them [6, 14]. The blood pressure (BP) decreased transiently during the increased respiratory movement and then rose again and maintained the original level for a while. The heart rate slowed slightly just after the gas mixture inhalation. It is be-

lieved that the initial BP depression with the decreased heart rate is due to the direct depressant effect of CO₂ upon the heart and that the return of blood pressure to its original level was due to the rapid rise of PaCO₂. After 30–90 min, the respiratory center was depressed, and hypotension occurred gradually or suddenly from reduced cardiac output, leading to an apnea and eventually to circulatory arrest [6].

In higher concentrations of CO₂, unconsciousness occurred almost instantaneously and respiratory movement ceased in 1 min. After a few minutes of apnea, circulatory arrest was seen. These findings show that the cause of death in breathing high concentrations of CO₂ is not the hypoxia but the intoxication of carbon dioxide [6].

CO₂ toxicity in humans Carbon dioxide at low concentration has little, if any, toxicological effects. At higher concentrations (>5%), it causes the development of hypercapnia and respiratory acidosis. Severe acidosis increases the effects of parasympathetic nervous activity, possibly by interfering the hydrolysis of acetylcholine by acetylcholinesterase, resulting in a depression of the respiration and the circulation [6]. Concentrations of more than 10% carbon dioxide may cause convulsions, coma, and death [1, 15]. CO₂ levels of more than 30% act rapidly leading to loss of consciousness in seconds. This would explain why victims of accidental intoxications often do not act to resolve the situation (open a door, etc.) [7, 10, 16].

Studies have shown a wide variability of CO₂ tolerance. Blood concentrations ranged between at least 0.055 and 0.085 atm. (41.8–64.6 mmHg) among subjects with symptoms, suggesting that a safe CO₂ exposure level cannot be characterized by a single value [10]. Concentrations of fatal cases of carbon dioxide vary between 14.1 and 26% CO₂ and an accompanying O₂ level between 4.2 and 25% [1, 8, 11]. It was also determined that CO₂ tolerance decreases with age ($p < 0.0001$) and suggested that smokers might have more tolerance due to habituation of higher CO₂ levels in cigarette smoke [10, 16].

Effects of oxygen treatment have been studied on animal models, and both normal and high concentration oxygen have been recommended in the literature [16, 17]. Niu et al. showed in a study performed on Sprague Dawley rats who inhaled carbon dioxide gas that the levels of serum troponin I (CTNI), CK, serum potassium (K), and AST are lower afterwards when treated with hyperbaric oxygen therapy compared to other oxygen treatments ($p < 0.05$). Levels of serum sodium (Na) and Chloride (Cl) were higher in hyperbaric therapy ($p < 0.05$). There was no significant difference in pH, PO₂, and PCO₂ among all oxygen-therapy groups ($p > 0.05$); however, there were significantly less pathological changes in the lungs with hyperbaric therapy [17]. On the other hand, high concentrations of oxygen raise venous pO₂ which reduces the solubility of carbon dioxide in the blood. With no changes in the metabolic conditions, this in turn causes pCO₂ in the venous blood to rise. Due to this so-called Haldane effect, an initial increase of pCO₂ in the bloodstream is to be expected when giving oxygen to a hypoxic carbon dioxide intoxicated person. It has also been suggested that due to high concentrations of oxygen, an increase in dead space is to be expected. P_{ET}CO₂ might therefore underestimate PaCO₂. Other studies have found that there is no significant effect between administering normal levels of oxygen or hyperoxic gas [16, 17].

Diagnosing CO₂ intoxication Diagnosing CO₂ intoxication is in the first place based on scene investigation and circumstances surrounding a victim [18]. Blood analysis will show increased carbon dioxide levels but is not always available pre-hospital [15]. Ischemic ECG changes (ST-segment changes and T-wave inversion) have also been described in several cases [19]. Victims might have burn wounds from contact with dry ice [15]. During massive geothermal emission events in Cameroon, cutaneous erythema and bullae have also been associated with coma states caused by exposure to carbon dioxide in the air on an unknown proportion of corpses and 161 (19%) survivors treated in a hospital. Though, it needs to be said that these were not reported in any other carbon dioxide case without contact to other acidic gasses [13].

Post-mortem diagnosis of carbon dioxide poisoning Post-mortem identification of carbon dioxide intoxication might even be more difficult. External examination of the body is often unremarkable [1, 7]. Although, bruises and hematoma might be present from CPR or from other rescue efforts [10]. Blood analysis for CO₂ content has only a limited diagnostic value, as CO₂ rapidly accumulates after death [18]. Nevertheless, a case study in Japan showed higher levels of CO₂ than those in the blood of deceased healthy persons. The levels found were similar to samples of fire victims, though there was no carbon monoxide (CO) found in this case (something that is usually found in fire victims) [8]. Analysis of lung gasses might also assist in determining the cause of death: an elevated CO₂ content in lung gasses has been described after vacuum degassing of the lungs in an argon atmosphere and quadrupole-mass-spectrometry [10]. Victims might also show passive hyperemia of the internal organs, pleural petechiae, and edema with a moderate congestion of the lungs and the brains [1, 7, 10]. Due to the limited possibilities of proving lethal CO₂ intoxications post-mortem, a close communication is needed among all of the involved parties, especially pre-hospital responders [9].

Management When confronted with patients with carbon dioxide intoxication, the immediate removal of the casualty (and unprotected rescuers) from the toxic environment is needed [15]. Oxygen should be administered, and appropriate supportive care is advisable [15]. Relating to treatment, both normal and high concentration oxygen have been recommended in the literature, without a definite consensus being reached at the moment. Benefits of each approach have been discussed earlier within the topic "CO₂ toxicity in humans" [16, 17].

Conclusions

Carbon dioxide poisoning, having a role both as an asphyxiant and as a toxicant, is a rare but not to miss diagnosis. Special attention is needed for pre-hospital responders, who should stay alert for the possibility of a CO₂ intoxication for their own safety, especially in cases involving dry ice or confined spaces. When confronted with a victim, he/she should be removed from the dangerous area as fast as possible and oxygen should be administered. Without adequate treatment, victims may show acute reduced cognitive performance, respiratory failure, and circulatory arrest.

Funding

None to declare

Authors' contributions

KP determined the concept of the article and design, was responsible for a major part of the acquisition of articles included in the review (article search), revised the article critically for important intellectual content, and assisted in revising the article. SV drafted the article, revised the article critically for important intellectual content, and assisted in revising the article. SS assisted in (re-)drafting the article to journal guidelines, revised the article critically for important intellectual content, and assisted in revising the article. CS assisted in the determination of concept and design, revised the article critically for important intellectual content, and assisted in revising the article. All authors read and approved the final manuscript.

Authors' information

Kris Permentier is an anaesthesiologist, emergency physician, and head of the ER-department AZ St. Blasius, Dendermonde, Belgium, and came as a pre-hospital doctor in contact with a carbon dioxide poisoning case with three patients.

Steven Vercammen is a co-assistant emergency medicine at the Ghent University, Belgium, and a MSc in disaster medicine and management at the Philadelphia University, USA.

Sylvia Soetaert is a co-assistant emergency medicine at the Ghent University, Belgium.

Christian Schellemans is a cardiologist and emergency physician at AZ St. Blasius, Dendermonde, Belgium.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Not applicable

Ethics approval and consent to participate

Not applicable

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Contributor Information

Kris Permentier, Email: kris.permentier@azsintblasius.be.

Steven Vercammen, Email: steven.vercammen@ugent.be.

Sylvia Soetaert, Email: Sylvia.soetaert@ugent.be.

Christian Schellemans, Email: Christian.schellemans@azsintblasius.be.

References

1. Zaba C, Marcinkowski JT, Wojtyla A, Tezyk A, Tobolski J, Zaba Z. Acute collective gas poisoning at work in a manure storage tank. *Ann Agric Environ Med*. 2011;18(2):448–451. [[PubMed](#)] [[Google Scholar](#)]
2. Romeo L, Prigioni P, Marcheselli S, Marchiori L, Cerpelloni M, Fiorini C, et al. Acute poisoning with carbon dioxide: report of 2 fatal cases. *Med Lav*. 2002;93:26–33. [[PubMed](#)] [[Google Scholar](#)]
3. Zugibe FT, Costello JT, Breithaupt MK, Zappi E, Allyn B. The confined space-hypoxia syndrome. *J Forensic Sci*. 1987;32:554–60. [[PubMed](#)] [[Google Scholar](#)]
4. National Traumatic Occupational Fatality database 1980-1989. *The National Institute for Occupational Safety and Health*. West Virginia: US Department of Health and Human Services; 1992. [[Google Scholar](#)]
5. New OSHA rescue requirements for confined space retrieval. Occupational Safety and Health Administration, USA. 2015. [[PubMed](#)]
6. Ikeda N, Takahashi H, Umetsu K, Suzuki T. The course of respiration and circulation in death by carbon dioxide poisoning. *Forensic Sci Int*. 1989;41(1):93–99. doi: 10.1016/0379-0738(89)90240-5. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
7. Srisont S, Chirachariyavej T, Peonim AVM. A carbon dioxide fatality from dry ice. *J Forensic Sci*. 2009;54(4):961–962. doi: 10.1111/j.1556-4029.2009.01057.x. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
8. Norimine E, Ishizawa F, Honda K, Uemura S. Suicide case of carbon dioxide poisoning using dry ice. *Jpn J Toxicol*. 2009;22(2):121–124. [[PubMed](#)] [[Google Scholar](#)]
9. Rupp WR, Thierauf A, Nadjem H, Vogt S. Suicide by carbon dioxide. *Forensic Sci Int*. 2013;231(1):e30–e32. doi: 10.1016/j.forsciint.2013.05.013. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
10. Kettner M, Ramsthaler F, Juhnke C, Bux R, Schmidt P. A fatal case of CO₂ intoxication in a fermentation tank. *J Forensic Sci*. 2013;58(2):556–558. doi: 10.1111/1556-4029.12058. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
11. Marcinkowski T, Koska W. Przypadki zatrucia dwutlenkiem węgla w silosach (cases of carbon dioxide poisoning in silos) *Med Prac*. 1968;19(1):117–119. [[Google Scholar](#)]
12. Sigurdsson H, Devine JD, Tchua FM, Presser FM, Pringle MKW, Evans WC. Origin of the lethal gas burst from Lake Monoun, Cameroun. *J Volcanol Geoth Res*. 1987;31(1–2):1–16. doi: 10.1016/0377-0273(87)90002-3. [[CrossRef](#)] [[Google Scholar](#)]

13. Baxter PJ, Kapila M, Mfonfu D. Lake Nyos disaster, Cameroon, 1986: the medical effects of large scale emission of carbon dioxide. *BMJ*. 1989;298(6685):1437–41. doi: 10.1136/bmj.298.6685.1437. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
14. Hill L, Flack M. The effect of excess of carbon dioxide and of want of oxygen upon the respiration and the circulation. *J Physiol*. 1908;37(2):77–101. doi: 10.1113/jphysiol.1908.sp001258. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
15. Langford NJ. Carbon dioxide poisoning. *Toxicol Rev*. 2005;24(4):229–235. doi: 10.2165/00139709-200524040-00003. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
16. Gill M, Natoli MJ, Vacchiano C, Macleod DB, Ikeda K, Qin M, et al. Effects of elevated oxygen and carbon dioxide partial pressures on respiratory function and cognitive performance. *J Appl Physiol*. 2014;117(4):406–412. doi: 10.1152/jappphysiol.00995.2013. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
17. Spitz WU, Spitz DJ. Asphyxia. In: Spitz WU, editor. *Spitz and Fischer's medicolegal investigation of death*. Springfield: Charles C Thomas; 2006. pp. 783–845. [[Google Scholar](#)]
18. Halpern P, Raskin Y, Sorkine P, Oganezov A. Exposure to extremely high concentrations of carbon dioxide: a clinical description of a mass casualty incident. *Ann Emerg Med*. 2004;43(2):196–9. doi: 10.1016/j.annemergmed.2003.08.003. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
19. Niu YM, Hao FT, Xue CJ, Xia YJ, Zhou S, Lu QS, et al. Study on the effect of different oxygen therapies on rats with acute carbon dioxide poisoning. *Hum Exp Toxicol*. 2012;31(2):126–133. doi: 10.1177/0960327111412806. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]